

Anti-DARPP-32 Antibody

Our Anti-DARPP-32 rabbit polyclonal primary antibody from PhosphoSolutions is produced in-house. It
Catalog # AN1353

Product Information

Application	WB, IHC
Primary Accession	Q6J410
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	22913


Additional Information

Gene ID	360616
Other Names	DARPP32 antibody, DARPP-32 antibody, Dopamine and cAMP regulated neuronal phosphoprotein 32 antibody, Dopamine and cAMP regulated neuronal phosphoprotein antibody, Dopamine and cAMP regulated phosphoprotein antibody, Dopamine and cAMP regulated phosphoprotein DARPP 32 antibody, Dopamine and cAMP regulated phosphoprotein DARPP32 antibody, Dopamine- and cAMP-regulated neuronal phosphoprotein antibody, FLJ20940 antibody, IPPD antibody, Neuronal phosphoprotein DARPP 32 antibody, PPP1R1B antibody, PPR1B_HUMAN antibody, Protein phosphatase 1 regulatory (inhibitor) subunit 1B antibody, Protein phosphatase 1 regulatory inhibitor subunit 1B antibody, Protein phosphatase 1 regulatory subunit 1B antibody
Target/Specificity	DARPP-32 is a dopamine (DA) and cAMP-regulated ~32 kDa phosphoprotein that is associated with dopaminoceptive neurons (Fienberg et al., 1998). The protein inhibits protein phosphatase I when it is phosphorylated on Thr-34. In contrast, when DARPP-32 is phosphorylated on Thr-75 the protein acts as an inhibitor of PKA (Bibb et al., 1999). Phosphorylation of DARPP-32 is thought to play a critical role in the regulation of dopaminergic neurotransmission. In addition, the activity of DARPP-32 is also thought to play important roles in the actions of alcohol, caffeine and Prozac® (Maldve et al., 2002; Lindskog et al., 2002; Svenningsson et al., 2002).
Dilution	WB~~1:1000 IHC~~1:100~500
Format	Antigen Affinity Purified from Pooled Serum
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-DARPP-32 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


Background

DARPP-32 is a dopamine (DA) and cAMP-regulated ~32 kDa phosphoprotein that is associated with dopaminergic neurons (Fienberg et al., 1998). The protein inhibits protein phosphatase I when it is phosphorylated on Thr-34. In contrast, when DARPP-32 is phosphorylated on Thr-75 the protein acts as an inhibitor of PKA (Bibb et al., 1999). Phosphorylation of DARPP-32 is thought to play a critical role in the regulation of dopaminergic neurotransmission. In addition, the activity of DARPP-32 is also thought to play important roles in the actions of alcohol, caffeine and Prozac® (Maldve et al., 2002; Lindskog et al., 2002; Svenningsson et al., 2002).

Images



Immunofluorescence staining of cultured CD1 mouse striatal cells showing specific labeling of DARPP-32 (cat. AN1353, green, 1:250) and Tuj1 (red). Cells and photo courtesy of QBM CellScience.



Western blot of rat hippocampal lysate showing specific immunolabeling of the ~32 kDa DARPP protein.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.