

Anti-Dynamin (Ser778) Antibody

Our Anti-Dynamin (Ser778) sheep polyclonal phosphospecific primary antibody from PhosphoSolutions is
Catalog # AN1371

Product Information

Application	WB
Primary Accession	P21575
Host	Sheep
Clonality	Polyclonal
Isotype	IgG
Calculated MW	97295


Additional Information

Gene ID	140694
Other Names	B dynamin antibody, D100 antibody, DNM 1 antibody, DNM antibody, DNM1 antibody, DYN1_HUMAN antibody, Dynamin antibody, Dynamin-1 antibody, Dynamin1 antibody
Target/Specificity	Dynamin is a member of a group of nerve terminal proteins called dephosphins that regulate synaptic vesicle endocytosis (Cousin et al., 2001; Graham et al., 2002; Tsuboi et al., 2002). Cyclin dependent protein kinase 5 phosphorylates dynamin at Ser-774 and Ser-778 that are the phosphorylation sites on dynamin phosphorylated in vivo (Tan et al., 2003). Phosphorylation of these sites on dynamin is thought to play a key role in synaptic vesicle trafficking.
Dilution	WB~~1:1000
Format	Antigen Affinity Purified from Pooled Serum
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-Dynamin (Ser778) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

Dynamin is a member of a group of nerve terminal proteins called dephosphins that regulate synaptic vesicle endocytosis (Cousin et al., 2001; Graham et al., 2002; Tsuboi et al., 2002). Cyclin dependent protein kinase 5 phosphorylates dynamin at Ser-774 and Ser-778 that are the phosphorylation sites on dynamin phosphorylated in vivo (Tan et al., 2003). Phosphorylation of these sites on dynamin is thought to play a key role in synaptic vesicle trafficking.

Images



Western blot of rat hippocampal lysate stimulated with forskolin showing specific immunolabeling of the ~95 kDa dynamin phosphorylated at Ser778 in the first lane (-). Phosphospecificity is shown in the second lane (+) where immunolabeling is completely eliminated by blot treatment with lambda phosphatase (λ-Ptase, 1200 units for 30 min).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.