

Anti-Elk-1 (Ser383) Antibody

Our Anti-Elk-1 (Ser383) rabbit polyclonal phosphospecific primary antibody from PhosphoSolutions is
Catalog # AN1374

Product Information

Application	WB
Primary Accession	P19419
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	44888

Additional Information


Gene ID	2002
Other Names	ELK 1 antibody, Elk1 antibody, ELK1 member of ETS oncogene family antibody, ELK1 protein antibody, ELK1, ETS transcription factor antibody, ELK1_HUMAN antibody, ELK2 member of ETS oncogene family antibody, ETS domain containing protein Elk 1 antibody, ETS domain containing protein Elk1 antibody, ETS domain protein Elk1 antibody, ETS domain-containing protein Elk-1 antibody, ETS like gene 1 antibody, Member of ETS oncogene family antibody, Oncogene Elk1 antibody, Tyrosine kinase (ELK1) oncogene antibody
Target/Specificity	Elk-1 is a transcription factor involved in mediating gene transcription in response to growth factors (Hill and Treisman, 1995). Elk-1 is thought to be phosphorylated by MAP kinase at Ser-383 and phosphorylation at this site is essential for the transcriptional activity of Elk-1 (Li et al., 2003). Phosphorylation of Elk-1 has also been implicated in synaptic plasticity in the adult hippocampus (Thiels et al., 2002).
Dilution	WB~~1:1000
Format	Antigen Affinity Purified from Pooled Serum
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-Elk-1 (Ser383) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

Elk-1 is a transcription factor involved in mediating gene transcription in response to growth factors (Hill and Treisman, 1995). Elk-1 is thought to be phosphorylated by MAP kinase at Ser-383 and phosphorylation at this

site is essential for the transcriptional activity of Elk-1 (Li et al., 2003). Phosphorylation of Elk-1 has also been implicated in synaptic plasticity in the adult hippocampus (Thiels et al., 2002).

Images



Western blot of recombinant Elk-1 showing specific immunolabeling of the ~46 kDa Elk-1 phosphorylated at Ser383 in the right lane (-). Phosphospecificity is shown in the left lane (+) where immunolabeling is blocked by preadsorption of the phosphopeptide used as the antigen, but not by the corresponding non-phosphopeptide (not shown).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.