10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Microtubule Associated Protein 2 C/D (MAP2C/D) Antibody

Our Anti-Microtubule Associated Protein 2 C/D (MAP2C/D) primary antibody from PhosphoSolutions is mo
Catalog # AN1438

Product Information

Application	WB, IHC, ICC
Primary Accession	P11137
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a
Clone Names	2C4
Calculated MW	199526

Additional Information

Gene ID	4133
Other Names	DKFZp686I2148 antibody, MAP 2 antibody, MAP dendrite specific antibody, MAP-2 antibody, MAP2 antibody, MAP2A antibody, MAP2B antibody, MAP2C antibody, Microtubule associated protein 2 antibody, Microtubule-associated protein 2 antibody, MTAP2_HUMAN antibody
Target/Specificity	Microtubules are 25nm diameter protein rods found in most kinds of eukaryotic cells. Microtubules are associated with a family of proteins called microtubule associated proteins (MAPs), which includes the protein τ (tau) and a group of proteins referred to as MAP1, MAP2, MAP3, MAP4 and MAP5 (Kindler & Gardner 1994). MAP2 is made up of two ~280 kDa bands referred to as MAP2a and MAP2b. A third lower molecular weight form, MAP2C and MAP2D, corresponds to a pair of protein bands running at ~70 kDa on SDS-PAGE gels. All these MAP2 forms are derived from a single gene by alternate transcription, and all share a C-terminal sequence which includes either three or four microtubule binding peptide sequences, which are very similar to those found in the related microtubule binding protein τ (tau). MAP2 isoforms are expressed only in neuronal cells and specifically in the perikarya and dendrites of these cells. MAP2C and MAP2D are expressed earlier in development than the MAP2a and MAP2b isoforms, so that this antibody is a more useful marker of neuronal development. MAP2 has been recently shown to be the specific receptor for the neurosteroid pregnenolone (FontaineLenore V. et al., 2006).
Dilution	WB~~1:1000 IHC~~1:100~500 ICC~~N/A
Format	Protein G Purified
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions	Anti-Microtubule Associated Protein 2 C/D (MAP2C/D) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

Microtubules are 25nm diameter protein rods found in most kinds of eukaryotic cells. Microtubules are associated with a family of proteins called microtubule associated proteins (MAPs), which includes the protein τ (tau) and a group of proteins referred to as MAP1, MAP2, MAP3, MAP4 and MAP5 (Kindler & Gardner 1994). MAP2 is made up of two ~280 kDa bands referred to as MAP2a and MAP2b. A third lower molecular weight form, MAP2C and MAP2D, corresponds to a pair of protein bands running at ~70 kDa on SDS-PAGE gels. All these MAP2 forms are derived from a single gene by alternate transcription, and all share a C-terminal sequence which includes either three or four microtubule binding peptide sequences, which are very similar to those found in the related microtubule binding protein τ (tau). MAP2 isoforms are expressed only in neuronal cells and specifically in the perikarya and dendrites of these cells. MAP2C and MAP2D are expressed earlier in development than the MAP2a and MAP2b isoforms, so that this antibody is a more useful marker of neuronal development. MAP2 has been recently shown to be the specific receptor for the neurosteroid pregnenolone (FontaineLenore V. et al., 2006).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.