

Anti-MeCP2 Antibody

Our Anti-MeCP2 primary antibody from PhosphoSolutions is mouse monoclonal. It detects human, mouse,
Catalog # AN1442

Product Information

Application	WB, IHC, ICC
Primary Accession	P51608
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Clone Names	5H12
Calculated MW	52441

Additional Information

Gene ID	4204
Other Names	AUTSX 3 antibody, AUTSX3 antibody, DKFZp686A24160 antibody, Mbd 5 antibody, Mbd5 antibody, MECP 2 antibody, MeCP 2 protein antibody, MeCP-2 protein antibody, Mecp2 antibody, MECP2_HUMAN antibody, Methyl CpG binding protein 2 (Rett syndrome) antibody, Methyl CpG binding protein 2 antibody, Methyl-CpG-binding protein 2 antibody, MRX 16 antibody, MRX 79 antibody, MRX16 antibody, MRX79 antibody, MRXS 13 antibody, MRXS13 antibody, MRXSL antibody, PPMX antibody, RS antibody, RTS antibody, RTT antibody, WBP 10 antibody, WBP10 antibody
Target/Specificity	MeCP2 (Methyl-CpG Binding Protein 2) is a chromosomal protein that binds to methylated DNA. It can bind specifically to a single methyl-CpG pair and is not influenced by sequences flanking the methyl-CpGs. MeCP2 has been shown to mediate transcriptional repression through interaction with histone deacetylase and the corepressor SIN3A (Nan et al., 1998). Defects in MeCP2 are the cause of Rett syndrome (RTT) (Amir et al., 1999). RTT is an X-linked dominant disease; it is a progressive neurologic developmental disorder and one of the most common causes of mental retardation in females.
Dilution	WB~~1:1000 IHC~~1:100~500 ICC~~N/A
Format	Protein G Purified
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-MeCP2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

MeCP2 (Methyl-CpG Binding Protein 2) is a chromosomal protein that binds to methylated DNA. It can bind specifically to a single methyl-CpG pair and is not influenced by sequences flanking the methyl-CpGs. MeCP2 has been shown to mediate transcriptional repression through interaction with histone deacetylase and the corepressor SIN3A (Nan et al., 1998). Defects in MeCP2 are the cause of Rett syndrome (RTT) (Amir et al., 1999). RTT is an X-linked dominant disease; it is a progressive neurologic developmental disorder and one of the most common causes of mental retardation in females.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.