

## Anti-MEF2C (Ser222) Antibody

Our Anti-MEF2C (Ser222) rabbit polyclonal phosphospecific primary antibody from PhosphoSolutions is  
Catalog # AN1446

### Product Information

---

Application	WB
Primary Accession	<a href="#">Q06413</a>
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	51221

### Additional Information

---

Gene ID	4208
Other Names	C5DELq14.3 antibody, DEL5q14.3 antibody, MADS box transcription enhancer factor 2 polypeptide C (myocyte enhancer factor 2C) antibody, MADS box transcription enhancer factor 2 polypeptide C antibody, MEF2C antibody, MEF2C_HUMAN antibody, Myocyte enhancer factor 2C antibody, Myocyte specific enhancer factor 2C antibody, Myocyte-specific enhancer factor 2C antibody, OTTHUMP00000222409 antibody, Similar to MADS box transcription enhancer factor 2 polypeptide C antibody
Target/Specificity	MEF2C, also known as MADS box transcription enhancer factor 2, polypeptide C, is one of 4 MEF2 (myocyte enhancer factor 2) transcription factors that encode proteins for development of skeletal muscle and brain proliferation and differentiation (McDermott et al, 1993), along with regulating stress-response during cardiac hypertrophy in mammals (Wu et al, 2015). Phosphorylation of MEF2C at serine 59 has been shown to be negatively regulated by integrin-linked kinase (ILK) (Dong et al, 2015). Recently, high MEF2C expression has been associated with a subset of acute myeloid leukemia (AML) patients with adverse-risk disease features and poor outcomes (Laszlo et al, 2015). Phosphorylation of MEF2C at serine 222 may play a key role in MEF2C signaling.
Dilution	WB~~1:1000
Format	Antigen Affinity Purified from Pooled Serum
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-MEF2C (Ser222) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

## Background

---

MEF2C, also known as MADS box transcription enhancer factor 2, polypeptide C, is one of 4 MEF2 (myocyte enhancer factor 2) transcription factors that encode proteins for development of skeletal muscle and brain proliferation and differentiation (McDermott et al, 1993), along with regulating stress-response during cardiac hypertrophy in mammals (Wu et al, 2015). Phosphorylation of MEF2C at serine 59 has been shown to be negatively regulated by integrin-linked kinase (ILK) (Dong et al, 2015). Recently, high MEF2C expression has been associated with a subset of acute myeloid leukemia (AML) patients with adverse-risk disease features and poor outcomes (Laszlo et al, 2015). Phosphorylation of MEF2C at serine 222 may play a key role in MEF2C signaling.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.