

Anti-Nestin Antibody

Our Anti-Nestin primary antibody from PhosphoSolutions is mouse monoclonal. It detects human, mouse,
Catalog # AN1463

Product Information

| | |
|--------------------------|------------------------|
| Application | WB, IHC, ICC |
| Primary Accession | P48681 |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype | IgG1 |
| Clone Names | 4D11 |
| Calculated MW | 177439 |

Additional Information

| | |
|---------------------------|--|
| Gene ID | 10763 |
| Other Names | ESTM 46 antibody, FLJ 21841 antibody, FLJ21841 antibody, Intermediate filament protein antibody, Nbla00170 antibody, nes antibody, NEST_HUMAN antibody, Nestin antibody |
| Target/Specificity | Nestin is a member of the class IV intermediate filament protein family which is expressed in neuronal stem cells (Lendahl et al., 1990). Nestin was originally identified as a result of the production of a series of monoclonal antibodies directed against epitopes expressed on formalin fixed embryo day 15 rat spinal cord (Hockfield et al., 1985)). One of these antibodies, called Rat 401, stained fibrous profiles in the developing nervous system, but not in the mature nervous system. By screening bacteriophage expression libraries with Rat 401, Lendahl et al. were able to isolate a cDNA encoding the protein to which Rat 401 antibody bound. The protein proved to be an unusual member of the intermediate filament family, containing an alpha-helical region homologous to that found in keratin and neurofilament subunits. |
| Dilution | WB~~1:1000 IHC~~1:100~500 ICC~~N/A |
| Format | Protein G Purified |
| Storage | Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | Anti-Nestin Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |
| Shipping | Blue Ice |

Background

Nestin is a member of the class IV intermediate filament protein family which is expressed in neuronal stem cells (Lendahl et al., 1990). Nestin was originally identified as a result of the production of a series of monoclonal antibodies directed against epitopes expressed on formalin fixed embryo day 15 rat spinal cord (Hockfield et al., 1985)). One of these antibodies, called Rat 401, stained fibrous profiles in the developing nervous system, but not in the mature nervous system. By screening bacteriophage expression libraries with Rat 401, Lendahl et al. were able to isolate a cDNA encoding the protein to which Rat 401 antibody bound. The protein proved to be an unusual member of the intermediate filament family, containing an alpha-helical region homologous to that found in keratin and neurofilament subunits.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.