

# Anti-Neurofilament NF-M Antibody

Our Anti-Neurofilament NF-M primary antibody from PhosphoSolutions is chicken polyclonal. It detects  
Catalog # AN1469

## Product Information

---

<b>Application</b>	WB, IHC, ICC
<b>Primary Accession</b>	<a href="#">P12839</a>
<b>Host</b>	Chicken
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgY
<b>Calculated MW</b>	95791

## Additional Information

---

<b>Gene ID</b>	24588
<b>Other Names</b>	150 kDa Neurofilament Medium antibody, 160 kDa neurofilament protein antibody, Medium polypeptide 150kDa antibody, NEF3 antibody, Neurofilament 3 antibody, Neurofilament 3 medium antibody, Neurofilament medium polypeptide antibody, Neurofilament protein M antibody, Neurofilament triplet M protein antibody, Neurofilament-3 (150 kD medium) antibody, NF-M antibody, NFM antibody
<b>Target/Specificity</b>	Neurofilaments are the 10 nm or intermediate filament proteins found specifically in neurons, and are composed predominantly of three major proteins called NF-L, NF-M and NF-H (Harris et al., 1991). NF-M is the neurofilament middle or medium molecular weight polypeptide and runs on SDS-PAGE gels at 145-160 kDa, with some variability across species boundaries. Antibodies to NF-M are useful for identifying neuronal cells and their processes in tissue sections and in tissue culture. NF-M antibodies can also be useful to visualize neurofilament accumulations seen in many neurological diseases, such as Amyotrophic Lateral Sclerosis (Lou Gehrig's disease) and Alzheimer's disease (Hu et al., 2002).
<b>Dilution</b>	WB~~1:1000 IHC~~1:100~500 ICC~~N/A
<b>Format</b>	Total IgY fraction
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Anti-Neurofilament NF-M Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
<b>Shipping</b>	Blue Ice

## Background

---

Neurofilaments are the 10 nm or intermediate filament proteins found specifically in neurons, and are composed predominantly of three major proteins called NF-L, NF-M and NF-H (Harris et al., 1991). NF-M is the neurofilament middle or medium molecular weight polypeptide and runs on SDS-PAGE gels at 145-160 kDa, with some variability across species boundaries. Antibodies to NF-M are useful for identifying neuronal cells and their processes in tissue sections and in tissue culture. NF-M antibodies can also be useful to visualize neurofilament accumulations seen in many neurological diseases, such as Amyotrophic Lateral Sclerosis (Lou Gehrig's disease) and Alzheimer's disease (Hu et al., 2002).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.