10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-NSF (N-ethylmaleimide sensitive fusion protein) Antibody

Our Anti-NSF (N-ethylmaleimide sensitive fusion protein) rabbit polyclonal primary antibody from Pho
Catalog # AN1499

Product Information

Application	WB
Primary Accession	Q9QUL6
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	82652

Additional Information

Gene ID	60355
Other Names	N ethylmaleimide sensitive factor antibody, N ethylmaleimide sensitive factor like protein antibody, N ethylmaleimide sensitive fusion protein antibody, N-ethylmaleimide-sensitive fusion protein antibody, NEM sensitive fusion protein antibody, NEM-sensitive fusion protein antibody, NSF antibody, NSF_HUMAN antibody, SKD 2 antibody, SKD2 antibody, Vesicle fusing ATPase antibody, Vesicle-fusing ATPase antibody, Vesicular fusion protein NSF antibody, Vesicular-fusion protein NSF antibody
Target/Specificity	NSF (N-ethylmaleimide sensitive fusion protein) is a critical component of the SNARE (soluble NSF attachment protein receptors) protein complex that is involved in synaptic vesicle trafficking. Specifically, NSF has been found to be essential in membrane fusion. Furthermore, NSF has been recently demonstrated to bind other protein complexes such as AMPA receptor subunits (GluR2), GATE-16, LMA-1 and Rabs suggesting a more diverse role in the assembly of various protein complexes (Whiteheart et al., 2004).
Dilution	WB~~1:1000
Format	Antigen Affinity Purified from Pooled Serum
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-NSF (N-ethylmaleimide sensitive fusion protein) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

NSF (N-ethylmaleimide sensitive fusion protein) is a critical component of the SNARE (soluble NSF attachment protein receptors) protein complex that is involved in synaptic vesicle trafficking. Specifically, NSF has been found to be essential in membrane fusion. Furthermore, NSF has been recently demonstrated to bind other protein complexes such as AMPA receptor subunits (GluR2), GATE-16, LMA-1 and Rabs suggesting a more diverse role in the assembly of various protein complexes (Whiteheart et al., 2004).