

## Anti-Syntaxin Antibody

Our Anti-Syntaxin primary antibody from PhosphoSolutions is mouse monoclonal. It detects human, rat, Catalog # AN1572

## **Product Information**

WB, IHC **Application Primary Accession Q16623** Reactivity Rat, Pig Host Mouse Clonality Monoclonal Isotype IgG1 **Clone Names** SP8 Calculated MW 33023

## **Additional Information**

Gene ID 6804

Other Names HPC 1 antibody, Neuron specific antigen HPC1 antibody, Neuron-specific

antigen HPC-1 antibody, OTTHUMP00000174615 antibody,

OTTHUMP00000174616 antibody, OTTHUMP00000174617 antibody, OTTHUMP00000174618 antibody, P35-1 antibody, STX1 antibody, STX1A antibody, STX1A\_HUMAN antibody, SYN1A antibody, Syntaxin 1A (brain)

antibody, Syntaxin 1A brain antibody, Syntaxin-1A antibody

**Target/Specificity** Syntaxins are a family of proteins involved in the docking of synaptic vesicles

with the plasma membrane. Syntaxin, along with synaptobrevin and SNAP-25, forms the SNARE complex which mediates synaptic vesicle fusion and exocytosis. Binding of Munc18-1 to the N-terminus of syntaxin 1 has been demonstrated to be essential for exocytic membrane fusion (Khvotchev et al., 2007). Decreased levels of phosphorylated syntaxin 1 in the prefrontal cortex

have recently been correlated with schizophrenia (Castillo et al., 2010).

**Dilution** WB~~1:1000 IHC~~1:100~500

Format Protein G Purified

**Storage** Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** Anti-Syntaxin Antibody is for research use only and not for use in diagnostic

or therapeutic procedures.

**Shipping** Blue Ice

## **Background**

Syntaxins are a family of proteins involved in the docking of synaptic vesicles with the plasma membrane. Syntaxin, along with synaptobrevin and SNAP-25, forms the SNARE complex which mediates synaptic vesicle fusion and exocytosis. Binding of Munc18-1 to the N-terminus of syntaxin 1 has been demonstrated to be essential for exocytic membrane fusion (Khvotchev et al., 2007). Decreased levels of phosphorylated syntaxin 1 in the prefrontal cortex have recently been correlated with schizophrenia (Castillo et al., 2010).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.