

Anti-Testicular Receptor 2 (TR2) Antibody

Our Anti-Testicular Receptor 2 (TR2) primary antibody from PhosphoSolutions is rabbit polyclonal. It

Catalog # AN1584

Product Information

Application	WB
Primary Accession	Q505F1
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	65476

Additional Information

Gene ID	22025
Other Names	NR2C 1 antibody, NR2C1 antibody, NR2C1_HUMAN antibody, NUCLEAR HORMONE RECEPTOR TR2 antibody, Nuclear receptor subfamily 2 Group C isoform antibody, Nuclear receptor subfamily 2 group C member 1 antibody, Orphan nuclear receptor TR2 antibody, Testicular receptor 2 antibody, TR 2 antibody, TR2 antibody, TR2-11 antibody
Target/Specificity	Testicular receptor 2 (TR2) is a member of the orphan nuclear receptor family. It is widely expressed at a low level throughout the adult testis. TR2 represses transcription and binds DNA directly interacting with HDAC3 and HDAC4 via DNA-binding domains (Franco et al., 2003). TR2 has also been implicated in regulation of estrogen receptor activity in mammary glands (Hun et al., 2002). In addition, TR2 has recently been shown to form a heterodimer with TR4 that can bind to the direct repeat 6 element of the hepatitis B virus (HBV) enhancer II region thus suppressing HBV gene expression (Lin et al., 2008).
Dilution	WB~~1:1000
Format	Neat Serum
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-Testicular Receptor 2 (TR2) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

Testicular receptor 2 (TR2) is a member of the orphan nuclear receptor family. It is widely expressed at a low level throughout the adult testis. TR2 represses transcription and binds DNA directly interacting with HDAC3

and HDAC4 via DNA-binding domains (Franco et al., 2003). TR2 has also been implicated in regulation of estrogen receptor activity in mammary glands (Hun et al., 2002). In addition, TR2 has recently been shown to form a heterodimer with TR4 that can bind to the direct repeat 6 element of the hepatitis B virus (HBV) enhancer II region thus suppressing HBV gene expression (Lin et al., 2008).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.