

# Anti-Tyrosine Hydroxylase Antibody

Our Anti-Tyrosine Hydroxylase rabbit polyclonal primary antibody from PhosphoSolutions is produced in house.  
Catalog # AN1597

## Product Information

<b>Application</b>	WB, IHC, ICC
<b>Primary Accession</b>	<a href="#">P04177</a>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Calculated MW</b>	55966

## Additional Information

<b>Gene ID</b>	25085
<b>Other Names</b>	Dystonia 14 antibody, DYT14 antibody, DYT5b antibody, EC 1.14.16.2 antibody, OTTHUMP00000011225 antibody, OTTHUMP00000011226 antibody, ple antibody, Protein Pale antibody, TH antibody, The antibody, TY3H_HUMAN antibody, TYH antibody, Tyrosine 3 hydroxylase antibody, Tyrosine 3 monooxygenase antibody, Tyrosine 3-hydroxylase antibody, Tyrosine 3-monooxygenase antibody, Tyrosine hydroxylase antibody
<b>Target/Specificity</b>	Tyrosine hydroxylase (TH) is the rate-limiting enzyme in the synthesis of the catecholamines dopamine and norepinephrine. TH antibodies can therefore be used as markers for dopaminergic and noradrenergic neurons in a variety of applications including depression, schizophrenia, Parkinson's disease and drug abuse (Kish et al., 2001; Zhu et al., 2000; Zhu et al., 1999). TH antibodies can also be used to explore basic mechanisms of dopamine and norepinephrine signaling (Witkovsky et al., 2000; Salvatore et al., 2001; Dunkley et al., 2004).
<b>Dilution</b>	WB~~1:1000 IHC~~1:100~500 ICC~~N/A
<b>Format</b>	Antigen Affinity Purified from Pooled Serum
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Anti-Tyrosine Hydroxylase Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
<b>Shipping</b>	Blue Ice

## Background

Tyrosine hydroxylase (TH) is the rate-limiting enzyme in the synthesis of the catecholamines dopamine and

norepinephrine. TH antibodies can therefore be used as markers for dopaminergic and noradrenergic neurons in a variety of applications including depression, schizophrenia, Parkinson's disease and drug abuse (Kish et al., 2001; Zhu et al., 2000; Zhu et al., 1999). TH antibodies can also be used to explore basic mechanisms of dopamine and norepinephrine signaling (Witkovsky et al., 2000; Salvatore et al., 2001; Dunkley et al., 2004).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.