

Anti-Tyrosine Hydroxylase (Ser31) Antibody

Our Anti-Tyrosine Hydroxylase (Ser31) rabbit polyclonal phosphospecific primary antibody from Phospho

Catalog # AN1599

Product Information

Application	WB, IHC, ICC
Primary Accession	P04177
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	55966

Additional Information

Gene ID	25085
Other Names	Dystonia 14 antibody, DYT14 antibody, DYT5b antibody, EC 1.14.16.2 antibody, OTTHUMP00000011225 antibody, OTTHUMP00000011226 antibody, ple antibody, Protein Pale antibody, TH antibody, The antibody, TY3H_HUMAN antibody, TYH antibody, Tyrosine 3 hydroxylase antibody, Tyrosine 3 monooxygenase antibody, Tyrosine 3-hydroxylase antibody, Tyrosine 3-monooxygenase antibody, Tyrosine hydroxylase antibody
Target/Specificity	Tyrosine hydroxylase (TH) is the rate-limiting enzyme in the synthesis of the catecholamines dopamine and norepinephrine. TH antibodies can therefore be used as markers for dopaminergic and noradrenergic neurons in a variety of applications including depression, schizophrenia, Parkinson's disease and drug abuse (Kish et al., 2001; Zhu et al., 2000; Zhu et al., 1999). TH antibodies can also be used to explore basic mechanisms of dopamine and norepinephrine signaling (Witkovsky et al., 2000; Salvatore et al., 2001; Dunkley et al., 2004). The activity of TH is also regulated by phosphorylation (Haycock et al., 1982; Haycock et al., 1992; Jedynak et al., 2002). Phospho-specific antibodies for the phosphorylation sites on TH can be used to great effect in studying this regulation and in identifying the cells in which TH phosphorylation occurs.
Dilution	WB~~1:1000 IHC~~1:100~500 ICC~~N/A
Format	Antigen Affinity Purified from Pooled Serum
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-Tyrosine Hydroxylase (Ser31) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

Tyrosine hydroxylase (TH) is the rate-limiting enzyme in the synthesis of the catecholamines dopamine and norepinephrine. TH antibodies can therefore be used as markers for dopaminergic and noradrenergic neurons in a variety of applications including depression, schizophrenia, Parkinson's disease and drug abuse (Kish et al., 2001; Zhu et al., 2000; Zhu et al., 1999). TH antibodies can also be used to explore basic mechanisms of dopamine and norepinephrine signaling (Witkovsky et al., 2000; Salvatore et al., 2001; Dunkley et al., 2004). The activity of TH is also regulated by phosphorylation (Haycock et al., 1982; Haycock et al., 1992; Jedynak et al., 2002). Phospho-specific antibodies for the phosphorylation sites on TH can be used to great effect in studying this regulation and in identifying the cells in which TH phosphorylation occurs.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.