



Anti-Ubiquitin C-terminal Hydrolase 1 (UCHL1) Antibody

Our Anti-Ubiquitin C-terminal Hydrolase 1 (UCHL1) primary antibody from PhosphoSolutions is mouse m Catalog # AN1601

Product Information

WB, IHC, ICC **Application Primary Accession** P09936 Reactivity Pig Host Mouse Clonality Monoclonal Isotype IgG1 **Clone Names** BH7 **Calculated MW** 24824

Additional Information

Gene ID Other Names

7345

Epididymis luminal protein 117 antibody, Epididymis secretory protein Li 53 antibody, HEL 117 antibody, HEL S 53 antibody, NDGOA antibody, Neuron cytoplasmic protein 9.5 antibody, OTTHUMP00000218137 antibody, OTTHUMP00000218139 antibody, OTTHUMP00000218140 antibody, OTTHUMP00000218141 antibody, Park 5 antibody, PARK5 antibody, PGP 9.5 antibody, PGP9.5 antibody, PGP9.5 antibody, PGP9.5 antibody, Ubiquitin C terminal esterase L1 antibody, Ubiquitin C terminal hydrolase antibody, Ubiquitin C terminal hydrolase L1 antibody, Ubiquitin carboxyl terminal hydrolase isozyme L1 antibody, Ubiquitin thioesterase L1 antibody, Ubiquitin thiolesterase antibody, Ubiquitin thiolesterase L1 antibody, UCH-L1 antibody, UCHL1 antibody, UCHL1 antibody, UCHL1 antibody

Target/Specificity

Ubiquitin C-terminal hydrolase 1 (UCHL1) is also known as ubiquitin carboxyl esterase L1, ubiquitin thiolesterase, neuron-specific protein PGP9.5 and Park5. It was originally identified as a major component of the neuronal cytoplasm from 2-dimensional gel analysis of brain tissues, and was given the name PGP9.5 (1). It was later found that ubiquitin C-terminal hydrolase enzyme activity was associated with the PGP9.5 protein (2). The ubiquitin C-terminal hydrolases cleave ubiquitin from other molecules. Regulation of the ubiquitin pathway is very important and many disease states are associated with defects in this pathway. Genetic knockout of UCHL1 in mice results in a motor neuron degeneration similar to the spontaneous gracile axonal dystrophy (gad) mutant mice (3). Point mutations in the UCHL1 gene are associated with some forms of human Parkinson's disease (4). Since UCHL1 is heavily expressed in neurons, it is released in large amounts following injury or degeneration, so the detection of UCHL1 in CSF and other bodily fluids can be used as a biomarker.

Dilution WB~~1:1000 IHC~~1:100~500 ICC~~N/A

Format Protein G Purified

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

PrecautionsAnti-Ubiquitin C-terminal Hydrolase 1 (UCHL1) Antibody is for research use

only and not for use in diagnostic or therapeutic procedures.

Shipping Blue Ice

Background

Ubiquitin C-terminal hydrolase 1 (UCHL1) is also known as ubiquitin carboxyl esterase L1, ubiquitin thiolesterase, neuron-specific protein PGP9.5 and Park5. It was originally identified as a major component of the neuronal cytoplasm from 2-dimensional gel analysis of brain tissues, and was given the name PGP9.5 (1). It was later found that ubiquitin C-terminal hydrolase enzyme activity was associated with the PGP9.5 protein (2). The ubiquitin C-terminal hydrolases cleave ubiquitin from other molecules. Regulation of the ubiquitin pathway is very important and many disease states are associated with defects in this pathway. Genetic knockout of UCHL1 in mice results in a motor neuron degeneration similar to the spontaneous gracile axonal dystrophy (gad) mutant mice (3). Point mutations in the UCHL1 gene are associated with some forms of human Parkinson's disease (4). Since UCHL1 is heavily expressed in neurons, it is released in large amounts following injury or degeneration, so the detection of UCHL1 in CSF and other bodily fluids can be used as a biomarker.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.