10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Visinin-Like Protein 1 (VSNL1) Antibody

Our Anti-Visinin-Like Protein 1 (VSNL1) mouse monoclonal primary antibody from PhosphoSolutions is p
Catalog # AN1608

Product Information

Application	WB, IHC, ICC
Primary Accession	P62760
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Clone Names	2D11
Calculated MW	22142

Additional Information

Gene ID	7447
Other Names	21 kDa CABP antibody, Hippocalcin like protein 3 antibody, Hippocalcin-like protein 3 antibody, HLP 3 antibody, HLP3 antibody, HPCAL 3 antibody, HPCAL3 antibody, HUVISL1 antibody, Neural visinin-like protein 1 antibody, Neural visinin-like type 1 protein antibody, Neurocalcin alpha antibody, NVL 1 antibody, NVL1 antibody, Nvp1 antibody, OZ1 antibody, Ratnvp1 antibody, VILIP antibody, VILIP1 antibody, VILIP-1 antibody, Visinin antibody, Visinin like 1 antibody, Visinin like protein 1 antibody, Visinin-like protein 1 antibody, VISL 1 antibody, VISL1 antibody, VISL1_HUMAN antibody, VLP-1 antibody, Vnsl1 antibody, Vsnl1 antibody
Target/Specificity	Visinin-like protein 1 (VSNL1), also known as VILIP1, is a calcium sensor protein expressed exclusively in neurons. Highest levels of VSNL1 expression are found in cerebellar Purkinje cells. VSNL1 has been implicated in the modulation of cell signaling cascades via regulation of adenyl cyclase activity (Braunewell et al., 1997). Additionally, VSNL1 has been associated with amyloid plaques and neurofibrillar tangles in Alzheimer's disease (Schnurra et al., 2001).
Dilution	WB~~1:1000 IHC~~1:100~500 ICC~~N/A
Format	Protein G Purified
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-Visinin-Like Protein 1 (VSNL1) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

Visinin-like protein 1 (VSNL1), also known as VILIP1, is a calcium sensor protein expressed exclusively in neurons. Highest levels of VSNL1 expression are found in cerebellar Purkinje cells. VSNL1 has been implicated in the modulation of cell signaling cascades via regulation of adenyl cyclase activity (Braunewell et al., 1997). Additionally, VSNL1 has been associated with amyloid plaques and neurofibrillar tangles in Alzheimer’s disease (Schnurra et al., 2001).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.