10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Arp3 (C-terminal region) Antibody

Catalog # AN1641

Product Information

Application	WB, IHC, ICC
Primary Accession	P61158
Reactivity	Rat
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	47371

Additional Information

Gene ID	10096
Other Names	ACTR3, Arp3
Target/Specificity	Cellular morphology, adhesion, and motility occur through dynamic reorganization of actin-based superstructures. Actin-binding proteins are critical for regulating actin polymerization and superstructure formation. The Arp2/3 complex is an actin polymerization-inducing complex that includes Arp2, Arp3, p41-Arc, p34-Arc, p21-Arc, p20-Arc, and p16-Arc. Several nucleation promoting factors, such as WASP and coronin, regulate the activity of the Arp2/3 complex. In addition, the Arp2/3 complex may be regulated by phosphorylation of specific subunits in the complex. Arp2 has two phosphosites, Thr-237 and Thr-238, that are evolutionarily conserved, and are phosphorylated along with Tyr-202 in response to growth factor stimulation. These phosphorylation events may regulate binding to the pointed end of actin filaments, and alanine substitutions of these Arp2 phosphosites inhibit membrane protrusions. Thus, phosphorylation may be another mode of Arp2/3 complex regulation in addition to the activity of nucleation-promoting factors.
Dilution	WB~~1:1000 IHC~~1:100~500 ICC~~N/A
Format	Antigen Affinity Purified
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-Arp3 (C-terminal region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

Cellular morphology, adhesion, and motility occur through dynamic reorganization of actin-based

superstructures. Actin-binding proteins are critical for regulating actin polymerization and superstructure formation. The Arp2/3 complex is an actin polymerization-inducing complex that includes Arp2, Arp3, p41-Arc, p34-Arc, p21-Arc, p20-Arc, and p16-Arc. Several nucleation promoting factors, such as WASP and coronin, regulate the activity of the Arp2/3 complex. In addition, the Arp2/3 complex may be regulated by phosphorylation of specific subunits in the complex. Arp2 has two phosphosites, Thr-237 and Thr-238, that are evolutionarily conserved, and are phosphorylated along with Tyr-202 in response to growth factor stimulation. These phosphorylation events may regulate binding to the pointed end of actin filaments, and alanine substitutions of these Arp2 phosphosites inhibit membrane protrusions. Thus, phosphorylation may be another mode of Arp2/3 complex regulation in addition to the activity of nucleation-promoting factors.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.