

# Anti-ATM (Ser-1981), Phosphospecific Antibody

Catalog # AN1647

## Product Information

---

|                          |                        |
|--------------------------|------------------------|
| <b>Application</b>       | WB, ICC, IP            |
| <b>Primary Accession</b> | <a href="#">Q13315</a> |
| <b>Host</b>              | Mouse                  |
| <b>Clonality</b>         | Mouse Monoclonal       |
| <b>Isotype</b>           | IgG1                   |
| <b>Clone Names</b>       | M366                   |
| <b>Calculated MW</b>     | 350687                 |

## Additional Information

---

|                    |  |
|--------------------|--|
| <b>Gene ID</b>     | 472  |
| <b>Other Names</b> | ataxia telangiectasia mutated, AT1 ATDC TEL1 TELO1 |

|                           |   |
|---------------------------|---|
| <b>Target/Specificity</b> | Ataxia telangiectasia mutated kinase (ATM) is a serine/threonine kinase that regulates cell cycle checkpoints and DNA repair. Mutations of ATM cause a spectrum of defects ranging from neurodegeneration to cancer predisposition. Activation of ATM after DNA damage involves Cdk5 mediated phosphorylation of Ser-794 followed by autophosphorylation at Ser-1891. Active ATM kinase regulates a number of proteins involved in cell cycle checkpoint control, apoptosis and DNA repair. The Cdk5-ATM pathway regulates phosphorylation and function of the ATM targets p53 and H2AX in postmitotic neurons. Other known substrates of ATM include Chk2, Chk1, CtIP, 4E-BP1, BRCA1, RPA3, SMC1, FANCD2, Rad17, Artemis, Nbs1, and the I-2 regulatory subunit of PP1. Thus, activation of Cdk5 by DNA damage may be an important initiator of ATM-dependent regulation of cell cycle checkpoints. |
|---------------------------|---|

|                 |                             |
|-----------------|-----------------------------|
| <b>Dilution</b> | WB~~1:1000 ICC~~N/A IP~~N/A |
|-----------------|-----------------------------|

|               |                    |
|---------------|--------------------|
| <b>Format</b> | Protein A Purified |
|---------------|--------------------|

|                |  |
|----------------|--|
| <b>Storage</b> | Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
|----------------|--|

|                    |   |
|--------------------|---|
| <b>Precautions</b> | Anti-ATM (Ser-1981), Phosphospecific Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |
|--------------------|---|

|                 |          |
|-----------------|----------|
| <b>Shipping</b> | Blue Ice |
|-----------------|----------|

## Background

---

Ataxia telangiectasia mutated kinase (ATM) is a serine/threonine kinase that regulates cell cycle checkpoints and DNA repair. Mutations of ATM cause a spectrum of defects ranging from neurodegeneration to cancer predisposition. Activation of ATM after DNA damage involves Cdk5 mediated phosphorylation of Ser-794

followed by autophosphorylation at Ser-1891. Active ATM kinase regulates a number of proteins involved in cell cycle checkpoint control, apoptosis and DNA repair. The Cdk5-ATM pathway regulates phosphorylation and function of the ATM targets p53 and H2AX in postmitotic neurons. Other known substrates of ATM include Chk2, Chk1, CtIP, 4E-BP1, BRCA1, RPA3, SMC1, FANCD2, Rad17, Artemis, Nbs1, and the I-2 regulatory subunit of PP1. Thus, activation of Cdk5 by DNA damage may be an important initiator of ATM-dependent regulation of cell cycle checkpoints.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.