

Anti-Caspase-3 (N-terminal region) Antibody

Catalog # AN1668

Product Information

Application	WB, ICC
Primary Accession	P42574
Host	Mouse
Clonality	Mouse Monoclonal
Isotype	IgG2a
Clone Names	M377
Calculated MW	31608

Additional Information

Gene ID	836
Other Names	Caspase, -3, CPP-32, Apoptain, Yama, SCA-1, CASP-2, NEDD2, ICH-1, p18, p13, p12
Target/Specificity	The caspases are a group of cysteine enzymes, which cleave proteins in response to intrinsic and extrinsic pathways that cause apoptotic cell death. The caspases can be grouped into two subgroups based on their roles in apoptosis. Initiator caspases (caspases 2, 8, 9, and 10) are activated through the apoptosis-signaling pathways and activate the effector caspases (caspases 3, 6, and 7) which carry out apoptosis. Caspase cascades are initiated through assembly of multiprotein complexes that trigger activation of the initiator caspases, which are then released and are able to activate the downstream effector caspases.
Dilution	WB~~1:1000 ICC~~N/A
Format	Protein A Purified
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-Caspase-3 (N-terminal region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

The caspases are a group of cysteine enzymes, which cleave proteins in response to intrinsic and extrinsic pathways that cause apoptotic cell death. The caspases can be grouped into two subgroups based on their roles in apoptosis. Initiator caspases (caspases 2, 8, 9, and 10) are activated through the apoptosis-signaling pathways and activate the effector caspases (caspases 3, 6, and 7) which carry out apoptosis. Caspase cascades are initiated through assembly of multiprotein complexes that trigger activation of the initiator

caspases, which are then released and are able to activate the downstream effector caspases.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.