

Anti-Caspase-3 (p17 subunit) Antibody

Catalog # AN1669

Product Information

| | |
|--------------------------|------------------------|
| Application | WB, IHC |
| Primary Accession | P42574 |
| Host | Mouse |
| Clonality | Mouse Monoclonal |
| Isotype | IgG1 |
| Clone Names | 31A1067 |
| Calculated MW | 31608 |

Additional Information

| | |
|--------------------|-----------------------------|
| Gene ID | 836 |
| Other Names | Yama, CPP32, apopain, CASP3 |

| | |
|---------------------------|--|
| Target/Specificity | The caspases are a group of cysteine enzymes, which cleave proteins in response to intrinsic and extrinsic pathways that cause apoptotic cell death. The caspases can be grouped into two subgroups based on their roles in apoptosis. Initiator caspases (caspases 2, 8, 9, and 10) are activated through the apoptosis-signaling pathways and activate the effector caspases (caspases 3, 6, and 7) which carry out apoptosis. Caspase cascades are initiated through assembly of multiprotein complexes that trigger activation of the initiator caspases, which are then released and are able to activate the downstream effector caspases. |
|---------------------------|--|

| | |
|-----------------|---------------------------|
| Dilution | WB~~1:1000 IHC~~1:100~500 |
|-----------------|---------------------------|

| | |
|---------------|--------------------|
| Format | Protein G Purified |
|---------------|--------------------|

| | |
|----------------|--|
| Storage | Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
|----------------|--|

| | |
|--------------------|---|
| Precautions | Anti-Caspase-3 (p17 subunit) Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |
|--------------------|---|

| | |
|-----------------|----------|
| Shipping | Blue Ice |
|-----------------|----------|

Background

The caspases are a group of cysteine enzymes, which cleave proteins in response to intrinsic and extrinsic pathways that cause apoptotic cell death. The caspases can be grouped into two subgroups based on their roles in apoptosis. Initiator caspases (caspases 2, 8, 9, and 10) are activated through the apoptosis-signaling pathways and activate the effector caspases (caspases 3, 6, and 7) which carry out apoptosis. Caspase cascades are initiated through assembly of multiprotein complexes that trigger activation of the initiator caspases, which are then released and are able to activate the downstream effector caspases.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.