

Anti-Cdk1 (Tyr-15) [conserved site], Phosphospecific Antibody

Catalog # AN1714

Product Information

Application	WB
Primary Accession	P06493
Host	Mouse
Clonality	Mouse Monoclonal
Isotype	IgG1
Clone Names	M231
Calculated MW	34095

Additional Information

Gene ID	983
Other Names	Cdc2

Target/Specificity	Cyclin-dependent kinases (Cdks) are a family of serine/threonine kinases that require association with regulatory subunits known as cyclins for activation. In addition, post-translational phosphorylation and dephosphorylation events regulate Cdk activity. Phosphorylation of Thr-160 in the T loop by Cdk-activating kinase (CAK) is an obligatory step in kinase activation. By contrast, phosphorylation of the Thr-14 and Tyr-15 residues by the Wee1 family of dual specificity kinases is inhibitory for the Cdks, and dephosphorylation of these residues by the Cdc25 family of phosphatases coincides with Cdk activation. Alternatively, Cdk5 appears to require different mechanisms for activation. This Cdk is activated through association with specific activators, including p35, p39, and p67. Cdk5 is primarily activated in neuronal cells, and only c-Abl kinase, rather than Wee family members, have been shown to phosphorylate Tyr-15 to regulate its activity.
---------------------------	---

Dilution	WB~~1:1000
Format	Protein A Purified
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-Cdk1 (Tyr-15) [conserved site], Phosphospecific Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

Cyclin-dependent kinases (Cdks) are a family of serine/threonine kinases that require association with

regulatory subunits known as cyclins for activation. In addition, post-translational phosphorylation and dephosphorylation events regulate Cdk activity. Phosphorylation of Thr-160 in the T loop by Cdk-activating kinase (CAK) is an obligatory step in kinase activation. By contrast, phosphorylation of the Thr-14 and Tyr-15 residues by the Wee1 family of dual specificity kinases is inhibitory for the Cdks, and dephosphorylation of these residues by the Cdc25 family of phosphatases coincides with Cdk activation. Alternatively, Cdk5 appears to require different mechanisms for activation. This Cdk is activated through association with specific activators, including p35, p39, and p67. Cdk5 is primarily activated in neuronal cells, and only c-Abl kinase, rather than Wee family members, have been shown to phosphorylate Tyr-15 to regulate its activity.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.