

Anti-CREB (N-terminal region) Antibody

Catalog # AN1723

Product Information

Application	WB, ICC, IP
Primary Accession	P16220
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	35136

Additional Information

Gene ID	1385
Other Names	Cyclic AMP, CREB

Target/Specificity	CREB (cyclic AMP response element-binding protein) is a stimulus-induced transcription factor that plays pivotal roles in cell survival and proliferation. CREB is expressed in various tissues, and has important gene-regulating roles in the nervous system. The transactivation function of CREB is primarily regulated through Ser-133 phosphorylation by cAMP-dependent protein kinase A (PKA) and related kinases. CREB is phosphorylated at other sites in response to calcium influx and DNA damage. The DNA-damage responsive nuclear kinase, HIPK2, can phosphorylate Ser-271 but not Ser-133 in CREB, and this phosphorylation activates CREB transactivation function. Mutation of Ser-271 to Glu-271 potentiates the CREB transactivation function. Thus, phosphorylation of Ser-271 may be the mode of activation for CREB-dependent transcription in response to genotoxic stress.
---------------------------	--

Dilution	WB~~1:1000 ICC~~N/A IP~~N/A
-----------------	-----------------------------

Format	Antigen Affinity Purified
---------------	---------------------------

Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
----------------	--

Precautions	Anti-CREB (N-terminal region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
--------------------	--

Shipping	Blue Ice
-----------------	----------

Background

CREB (cyclic AMP response element-binding protein) is a stimulus-induced transcription factor that plays pivotal roles in cell survival and proliferation. CREB is expressed in various tissues, and has important gene-regulating roles in the nervous system. The transactivation function of CREB is primarily regulated through Ser-133 phosphorylation by cAMP-dependent protein kinase A (PKA) and related kinases. CREB is

phosphorylated at other sites in response to calcium influx and DNA damage. The DNA-damage responsive nuclear kinase, HIPK2, can phosphorylate Ser-271 but not Ser-133 in CREB, and this phosphorylation activates CREB transactivation function. Mutation of Ser-271 to Glu-271 potentiates the CREB transactivation function. Thus, phosphorylation of Ser-271 may be the mode of activation for CREB-dependent transcription in response to genotoxic stress.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.