

Anti-Crk II (Tyr-251), Phosphospecific Antibody

Catalog # AN1728

Product Information

| Application Primary Accession Reactivity Host Clonality Isotype Calculated MW | WB, IHC P46108 Rat Rabbit Rabbit Polyclonal IgG 33831 |
|---|--|
| | |
| Gene ID Other Names | 1398 v-crk sarcoma virus CT10 oncogene homolog, CRKII, CRKL |
| Target/Specificity | The Crk family of adaptor proteins (Crk I, Crk II and CrkL) are Src Homology 2 (SH2) and Src Homology 3 (SH3) domain-containing proteins that form protein complexes important for transmiting signals downstream of tyrosine kinases. Both Crk II and CrkL are composed of a single SH2 domain, followed by two tandem SH3 domains. Crk II is also alternatively spliced to a minor product, Crk I, which is structurally and functionally more similar to the v-Crk oncogene. Both Crk II and CrkL are ubiquitously expressed and their SH domains are highly homologous, however both are required for mouse development and have distinct non-overlapping phenotypes in knockout mice. Phosphorylation may be important for regulating Crk activity. Crk II Tyr-221 (CrkL Tyr-207) phosphorylation is a negative regulatory site, while Crk Tyr-251 phosphorylation in the SH3 domain is a positive regulatory site. EGF stimulation induces phosphorylation of Tyr-251, which increases binding of Crk to the SH2 domain of Abl, and promotes transactivation of Abl. |
| Dilution | WB~~1:1000 IHC~~1:100~500 |
| Format | Antigen Affinity Purified |
| Storage | Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | Anti-Crk II (Tyr-251), Phosphospecific Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |
| Shipping | Blue Ice |

Background

The Crk family of adaptor proteins (Crk I, Crk II and CrkL) are Src Homology 2 (SH2) and Src Homology 3 (SH3) domain-containing proteins that form protein complexes important for transmiting signals

downstream of tyrosine kinases. Both Crk II and CrkL are composed of a single SH2 domain, followed by two tandem SH3 domains. Crk II is also alternatively spliced to a minor product, Crk I, which is structurally and functionally more similar to the v-Crk oncogene. Both Crk II and CrkL are ubiquitously expressed and their SH domains are highly homologous, however both are required for mouse development and have distinct non-overlapping phenotypes in knockout mice. Phosphorylation may be important for regulating Crk activity. Crk II Tyr-221 (CrkL Tyr-207) phosphorylation is a negative regulatory site, while Crk Tyr-251 phosphorylation in the SH3 domain is a positive regulatory site. EGF stimulation induces phosphorylation of Tyr-251, which increases binding of Crk to the SH2 domain of Abl, and promotes transactivation of Abl.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.