

Anti-Cyclophilin B Antibody

Catalog # AN1738

Product Information

Application	WB, ICC, IP
Primary Accession	P23284
Host	Mouse
Clonality	Mouse Monoclonal
Isotype	IgG1
Clone Names	M019
Calculated MW	23743

Additional Information

Gene ID	5479
Other Names	Peptidyl-prolyl cis-trans isomerase B, PPIase B, CYP-S1, Cyclophilin B, Rotamase B, S-cyclophilin, SCYLP, PPIB, CYPB
Target/Specificity	Cyclophilins are a highly conserved family of peptidylprolyl cis-trans-isomerases (PPI) that are targets of the immunosuppressant drug cyclosporin A. These isomerases have been proposed to aid in protein folding. Cyclophilin B (PPIB) is an ER-localized chaperone protein that associates with type I procollagen, and facilitates procollagen sorting and transport through the secretory compartment. Mutations in the cyclophilin B gene lead to aberrant biosynthesis of type I procollagen, and osteogenesis imperfecta, a disorder characterized by bone fragility. Cyclophilin B has also been implicated as a marker in several cancers, including glioblastomas, neuroblastomas, and gastric cancers. In gastric cancer, Stat3 suppression of miR-520d-5p leads to upregulation of its downstream target cyclophilin B, and enables the growth and survival of these cancer cells.
Dilution	WB~~1:1000 ICC~~N/A IP~~N/A
Format	Protein G Purified
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-Cyclophilin B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

Cyclophilins are a highly conserved family of peptidylprolyl cis-trans-isomerases (PPI) that are targets of the immunosuppressant drug cyclosporin A. These isomerases have been proposed to aid in protein folding.

Cyclophilin B (PPIB) is an ER-localized chaperone protein that associates with type I procollagen, and facilitates procollagen sorting and transport through the secretory compartment. Mutations in the cyclophilin B gene lead to aberrant biosynthesis of type I procollagen, and osteogenesis imperfecta, a disorder characterized by bone fragility. Cyclophilin B has also been implicated as a marker in several cancers, including glioblastomas, neuroblastomas, and gastric cancers. In gastric cancer, Stat3 suppression of miR-520d-5p leads to upregulation of its downstream target cyclophilin B, and enables the growth and survival of these cancer cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.