

Anti-Estrogen Receptor α (C-terminus) Antibody

Catalog # AN1787

Product Information

Application	WB, ICC, IP
Primary Accession	P03372
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	66216

Additional Information

Gene ID	2099
Other Names	ESR, ESR1, ESRA, Estradiol receptor, Eralpha, ER

Target/Specificity	Estrogen receptor α (ER α) is a member of the steroid receptor superfamily and its structure includes an N-terminal ligand-independent transactivation domain (AF-1), a highly conserved DNA binding domain, and a C-terminal ligand-dependent transactivation domain (AF-2). AF-1 and AF-2 activate transcription independently and synergistically, and act in a promoter- and cell-specific manner. Phosphorylation at multiple sites provides an important mechanism to regulate ER α activity. Ser-104, Ser-106, Ser-118, and Ser-167 are located in the amino-terminal transcription activation function domain AF-1, and phosphorylation of these serine residues plays an important role in regulating ER α activity. In addition to these sites, phosphorylation of Tyr-537 has been implicated in maximal hormone binding, dimerization, and transcriptional activity. Tyr-537, located in the AF-2 domain, is phosphorylated by c-Src leading to nuclear export of ER α and degradation. Thus, a variety of phosphorylation events control ER α activity.
---------------------------	---

Dilution	WB~~1:1000 ICC~~N/A IP~~N/A
-----------------	-----------------------------

Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
----------------	--

Precautions	Anti-Estrogen Receptor α (C-terminus) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
--------------------	---


Shipping	Blue Ice
-----------------	----------

Background


Estrogen receptor α (ER α) is a member of the steroid receptor superfamily and its structure includes an N-terminal ligand-independent transactivation domain (AF-1), a highly conserved DNA binding domain, and a C-terminal ligand-dependent transactivation domain (AF-2). AF-1 and AF-2 activate transcription independently and synergistically, and act in a promoter- and cell-specific manner. Phosphorylation at

multiple sites provides an important mechanism to regulate ERα activity. Ser-104, Ser-106, Ser-118, and Ser-167 are located in the amino-terminal transcription activation function domain AF-1, and phosphorylation of these serine residues plays an important role in regulating ERα activity. In addition to these sites, phosphorylation of Tyr-537 has been implicated in maximal hormone binding, dimerization, and transcriptional activity. Tyr-537, located in the AF-2 domain, is phosphorylated by c-Src leading to nuclear export of ERα and degradation. Thus, a variety of phosphorylation events control ERα activity.

Images



Western blot image of human MCF-7 cells treated with pervanadate (1 mM) for 30 min. (lanes 1-6). Some lanes of the blot were then treated with alkaline phosphatase (lanes 2, 4, & 6). The blot was probed with mouse monoclonal anti-ERα (Tyr-537) phospho-specific (lanes 1 & 2), rabbit polyclonal anti-ERα (C-terminus) (lanes 3 & 4), and rabbit polyclonal anti-ERα (Tyr-537) phospho-specific (lanes 5 & 6).



Immunocytochemical labeling of Estrogen Receptor α in paraformaldehyde fixed and NP-40 permeabilized MDA-MB-231 cells. The cells were labeled with rabbit polyclonal anti-Estrogen Receptor α (AN1787). The antibody was detected using goat anti-rabbit DyLight® 594.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.