

Anti-Girdin (Tyr-1764), Phosphospecific Antibody

Catalog # AN1803

Product Information

Application	WB
Primary Accession	Q3V6T2
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	216042

Additional Information


Gene ID	55704
Other Names	APE, Galpha, vesicle, GIV, Girders actin filament, HkRP1, GRDN, CCDC88A
Target/Specificity	Girdin, a member of the CCDC88 (Hook related protein) family, is an actin binding protein involved with cell migration and maintaining cytoskeletal organization. Girdin has conserved domains at the N- and C-terminus that bind microtubules and actin, respectively. It enhances PI3-kinase dependent phosphorylation of proteins, most notably Akt. This same activity can contribute to tumor proliferation, invasion, and metastasis in breast, ovarian, prostate, and pancreatic tissues. Girdin is phosphorylated at three separate locations: Ser-1416, Ser-1674, and Tyr-1764. Ser-1416 is the primary Akt phosphorylation site, while Cyclin-dependent kinases interact with Girdin and phosphorylate Ser-1674. Multiple receptor tyrosine kinases can bind girdin and phosphorylate Tyr-1764.
Dilution	WB~~1:1000
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-Girdin (Tyr-1764), Phosphospecific Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

Girdin, a member of the CCDC88 (Hook related protein) family, is an actin binding protein involved with cell migration and maintaining cytoskeletal organization. Girdin has conserved domains at the N- and C-terminus that bind microtubules and actin, respectively. It enhances PI3-kinase dependent phosphorylation of proteins, most notably Akt. This same activity can contribute to tumor proliferation, invasion, and metastasis in breast, ovarian, prostate, and pancreatic tissues. Girdin is phosphorylated at three separate locations: Ser-1416, Ser-1674, and Tyr-1764. Ser-1416 is the primary Akt phosphorylation site, while Cyclin-dependent kinases interact with Girdin and phosphorylate Ser-1674. Multiple receptor tyrosine

kinases can bind girdin and phosphorylate Tyr-1764.

Images



Western blot image of human A431 cell lysates treated with pervanadate (lanes 1-4). The blot was treated with alkaline phosphatase to dephosphorylate Girdin phosphosites (lanes 2 & 4). The blot was probed with mouse monoclonal anti-Girdin (lanes 1 & 2) or rabbit polyclonal anti-Girdin (Tyr-1764), phospho-specific (lanes 3 & 4).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.