

Anti-GM130 (C-terminal region) Antibody

Catalog # AN1804

Product Information

Application	WB, ICC
Primary Accession	<u>Q08379</u>
Host	Mouse
Clonality	Mouse Monoclonal
Isotype	IgG1
Clone Names	M342
Calculated MW	113086

Additional Information

Gene ID Other Names	2801 GOLGA2, golgi autoantigen
Target/Specificity	Golgi Matrix Protein (GM130) is a peripheral cytoplasmic protein that is bound to Golgi membranes. It maintains cis-Golgi structure, and it regulates the disassembly and reassembly of the Golgi complex during mitosis. GM130 may also be important during docking and fusion of coatomer (COPI) coated vesicles to the Golgi membrane. The carboxy-terminal domain of GM130 is highly homologous to the human auto-antigen, golgin-95. GM130 interacts in a GTP-dependent manner with Rab1b protein, a regulator of anterograde traffic between ER and Golgi membranes. It has also been implicated in the activation of Ste-kinase, YSK1, during the golgi reorganization that occurs along with cell migration.
Dilution	WB~~1:1000 ICC~~N/A
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-GM130 (C-terminal region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

Golgi Matrix Protein (GM130) is a peripheral cytoplasmic protein that is bound to Golgi membranes. It maintains cis-Golgi structure, and it regulates the disassembly and reassembly of the Golgi complex during mitosis. GM130 may also be important during docking and fusion of coatomer (COPI) coated vesicles to the Golgi membrane. The carboxy-terminal domain of GM130 is highly homologous to the human auto-antigen, golgin-95. GM130 interacts in a GTP-dependent manner with Rab1b protein, a regulator of anterograde traffic between ER and Golgi membranes. It has also been implicated in the activation of Ste-kinase, YSK1, during the golgi reorganization that occurs along with cell migration.



Immunocytochemical labeling of GM130 in C2C12 cells. The cells were double-labeled with GM130 (golgi protein) mouse monoclonal and nSMase-3 rabbit polyclonal antibodies, then detected using appropriate secondary antibody conjugated to Cy2 or Cy3.



Western blot analysis of GM130 expression in human cells: MCF7 breast carcinoma (lane 1), A549 adenocarcinoma (lane 2), and MDA-MB-231 breast carcinoma (lane 3). The blot was probed with mouse monoclonal anti-GM130 (C-terminal region) at 1:500.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.