10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Anti-LIMK1 (C-terminus) Antibody

Catalog # AN1833

## Product Information

| | |
|---|---|
| **Application** | WB |
| **Primary Accession** | P53667 |
| **Host** | Rabbit |
| **Clonality** | Rabbit Polyclonal |
| **Isotype** | IgG |
| **Calculated MW** | 72585 |

## Additional Information


| | |
|---|---|
| **Gene ID** | 3984 |
| **Other Names** | LIMK |
| **Target/Specificity** | LIM kinases (LIMK1 and LIMK2) are serine/threonine kinases that have two zinc finger motifs, known as LIM motifs, in their amino-terminal regulatory domains. LIM kinases are involved in actin cytoskeletal regulation downstream of Rho-family GTPases, PAKs, and ROCK. PAK1 and ROCK phosphorylate LIMK1 or LIMK2 at the conserved Thr-508 or Thr-505 residues in the activation loop, increasing LIMK activity. In addition, VEGF-induced stress fiber formation has been linked to p38-mediated activation of LIMK through MK-2 phosphorylation of Ser-323. Activated LIM kinases inhibit the actin depolymerization activity of cofilin by phosphorylation at the amino-terminal Ser-3 residue of cofilin. In addition, LIMKs may have a function in the nucleus. It has been shown that the nuclear localization of LIMKs can mediate suppression of Rac/Cdc42-mediated cyclin D1 expression. This effect of LIMKs was independent of cofilin phosphorylation and the regulation of actin dynamics. |
| **Dilution** | WB~~1:1000 |
| **Storage** | Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| **Precautions** | Anti-LIMK1 (C-terminus) Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |
| **Shipping** | Blue Ice |

## Background

LIM kinases (LIMK1 and LIMK2) are serine/threonine kinases that have two zinc finger motifs, known as LIM motifs, in their amino-terminal regulatory domains. LIM kinases are involved in actin cytoskeletal regulation downstream of Rho-family GTPases, PAKs, and ROCK. PAK1 and ROCK phosphorylate LIMK1 or LIMK2 at the conserved Thr-508 or Thr-505 residues in the activation loop, increasing LIMK activity. In addition,

VEGF-induced stress fiber formation has been linked to p38-mediated activation of LIMK through MK-2 phosphorylation of Ser-323. Activated LIM kinases inhibit the actin depolymerization activity of cofilin by phosphorylation at the amino-terminal Ser-3 residue of cofilin. In addition, LIMKs may have a function in the nucleus. It has been shown that the nuclear localization of LIMKs can mediate suppression of Rac/Cdc42-mediated cyclin D1 expression. This effect of LIMKs was independent of cofilin phosphorylation and the regulation of actin dynamics.

## Images




Western blot image of activated mouse recombinant LIMK1 untreated (lanes 1 & 3) or treated with lambda phosphatase (lanes 2 & 4). The blots were probed with anti-LIMK1 (C-term.) (lanes 1 & 2) and anti-LIMK1 (Thr-508) (lanes 3 & 4).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.