10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-eNOS (Ser-1177), Phosphospecific Antibody

Catalog # AN1866

Product Information

Application	WB
Primary Accession	P29474
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	133275

Additional Information


Gene ID	4846
Other Names	endothelial Nitric Oxide Synthase, eNOS, ecNOS, NOS-III, NOS3, NOSIII
Dilution	WB~~1:1000
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-eNOS (Ser-1177), Phosphospecific Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

Nitric oxide (NO) has a broad range of biological activities and is implicated in signaling pathways in phylogenetically diverse species. Nitric oxide synthases (NOS), the enzymes responsible for synthesis of NO, are homodimers whose monomers are themselves two fused enzymes: a cytochrome reductase and a cytochrome that requires three cosubstrates (L-arginine, NADPH, and oxygen) and five cofactors or prosthetic groups (FAD, FMN, calmodulin, tetrahydrobiopterin, and heme). Several distinct NOS isoforms are produced from three distinct genes. The inducible form of NOS, iNOS (NOS-II), is Ca2+ independent and is expressed in a broad range of cell types, and two constitutive Ca2+/CaM-dependent forms of NOS: nNOS (bNOS, NOS-I) identified in neurons and eNOS (ecNOS, NOS-III) identified in endothelial cells. Regulation of eNOS activity occurs through phosphorylation at multiple sites. Phosphorylation of Ser-633 (mouse Ser-632) in the FMN binding domain increases eNOS activity and may be important for the maintenance of NO synthesis after initial activation by Ca2+ flux and Ser-1177 phosphorylation.

Images

Western blot analysis of human umbilical vein endothelial cells treated with calyculin A (100 nM) for 30 min. (lanes 1, 3, 5, & 7) then the blots were treated with lambda phosphatase (lanes 2, 4, 6, & 8). The blots were probed



with anti-endothelial nitric oxide synthase (eNOS) (C-terminal region) (lanes 1 & 2), anti-eNOS (Ser-632) (lanes 3 & 4), anti-eNOS (Ser-1177) (lanes 5 & 6), or anti-eNOS (a.a. 1172-1181) (lanes 7 & 8).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.