

# Anti-Paxillin (Ser-178), Phosphospecific Antibody

Catalog # AN1890

## Product Information

---

<b>Application</b>	WB, IHC
<b>Primary Accession</b>	<a href="#">P49023</a>
<b>Host</b>	Rabbit
<b>Clonality</b>	Rabbit Polyclonal
<b>Isotype</b>	IgG
<b>Calculated MW</b>	64505

## Additional Information

---

<b>Gene ID</b>	5829
<b>Other Names</b>	Paxillin, PXN
<b>Dilution</b>	WB~~1:1000 IHC~~1:100~500
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Anti-Paxillin (Ser-178), Phosphospecific Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
<b>Shipping</b>	Blue Ice

## Background


---

Paxillin, a focal adhesion protein, is involved in focal adhesion formation during cell adhesion and migration. Paxillin contains LD motifs, LIM domains, and SH3-/SH2-binding domains that participate in a variety of protein-protein interactions with kinases, GTPase-activating proteins, and cytoskeletal proteins. Phosphorylation of paxillin occurs at both tyrosine and serine sites. Serine phosphorylation of paxillin occurs in response to growth-factor activation and fibronectins. Both JNK1 and cdc2 kinases can phosphorylate serine 178 in paxillin. The mutant form of paxillin (S178A) decreases the migration of keratocytes and epithelial cells. Thus, phosphorylation paxillin at serine 178 may be important during cell migration

## Images

---

Western blot analysis of A431 cells (20 µg/lane) serum starved overnight and treated with EGF (100 ng/ml) for 5 min. The blot was probed with anti-Paxillin mouse monoclonal (PM1071) or anti-Paxillin (Ser-178) rabbit polyclonal (AN1890).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.