

Anti-Semaphorin-3A (C-terminal) Antibody

Catalog # AN1944

Product Information

| | |
|--------------------------|------------------------|
| Application | WB |
| Primary Accession | Q14563 |
| Host | Rabbit |
| Clonality | Rabbit Polyclonal |
| Isotype | IgG |
| Calculated MW | 88889 |

Additional Information


| | |
|--------------------|--|
| Gene ID | 10371 |
| Other Names | Sema3A, Semaphorin III |
| Dilution | WB~~1:1000 |
| Storage | Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | Anti-Semaphorin-3A (C-terminal) Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |
| Shipping | Blue Ice |

Background

One family of inhibitory axon guidance molecules is the semaphorins. The semaphorins include secreted, transmembrane, and GPI-anchored extracellular molecules that are involved in regulating axon guidance by inhibiting axons from growing toward incorrect targets. Semaphorin 3A (Sema3A) may play a particularly interesting role in limiting axon regeneration since it is expressed in meningeal fibroblasts that invade the injured spinal cord and surround the glial scar. In addition, the Sema3A co-receptors, Neuropilin-1 and Plexin-A1, are expressed on axons that regenerate up to the injured region, but do not cross this Sema3A-containing region. Thus, Sema3A and its co-receptors may have important roles in regulating axon guidance during neuronal development and after neuronal injury.

Images

Western blot image of human A431 cells and (lane 1) and human recombinant Sema3A/Fc chimera (95/125 kDa) (lane 2). The blots were probed with rabbit polyclonal anti-Semaphorin 3A (AN1944) at a dilution of 1:1000.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.