

# Anti-SHP2 (N-terminal region) Antibody

Catalog # AN1956

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">Q06124</a>
<b>Host</b>	Mouse
<b>Clonality</b>	Mouse Monoclonal
<b>Isotype</b>	IgG1
<b>Clone Names</b>	M163
<b>Calculated MW</b>	68011

## Additional Information

---

<b>Gene ID</b>	5781
<b>Other Names</b>	PTP1D, SHPTP2, Syp
<b>Dilution</b>	WB~~1:1000
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Anti-SHP2 (N-terminal region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
<b>Shipping</b>	Blue Ice

## Background


---

SHP2 (PTP1D, SH-PTP2, or Syp) is a widely expressed protein-tyrosine phosphatase (PTP) that maintains phosphotyrosine homeostasis during growth factor, cytokine, hormone and antigen receptor signaling. This phosphatase contains two N-terminal SH2 domains and a C-terminal phosphatase domain. SHP2 associates with EGF and PDGF growth factor receptors and is activated after stimulation of these receptors. Activation of SHP-2 and its association with Gab1 is critical for sustained ERK activation downstream of both growth factor and cytokine receptors. In addition to its role in Gab1-mediated Erk activation, SHP-2 attenuates EGF-dependent PI3 kinase activation by dephosphorylating Gab1 p85 binding sites. Thus, SHP2 is critical for maintaining phosphotyrosine homeostasis in many cell signaling pathways

## Images

---

Western blot analysis of adult mouse brain. The blot was probed with anti-SHP2 (N-terminal) antibody at 1:250 (lane 1), 1:500 (lane 2), 1:1000 (lane 3), and 1:2000 (lane 4).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.