

Anti-N-WASP Antibody

Catalog # AN2018

Product Information

Application	WB
Primary Accession	O00401
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	54827

Additional Information


Gene ID	8976
Other Names	Neural Wiskott-Aldrich syndrome protein, WASL, WASP
Dilution	WB~~1:1000
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-N-WASP Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

Members of the Wiskott-Aldrich syndrome protein (WASP) family regulate the formation of actin-based cell structures in many cell types. These proteins contain C-terminal actin-binding domains that can stimulate actin polymerization. In addition, these proteins bind the ARP2/3 complex, which can nucleate actin polymerization at sites that lead to branched actin structures. WASP is expressed primarily in hematopoietic cells, while its homolog N-WASP is widely expressed. These proteins have 48% identity in human with the highest homology in the functional regions of these proteins. Serine and tyrosine phosphorylation regulates the activity of both proteins. WASP is observed as a 63 kDa protein in hematopoietic cells, while N-WASP is observed as a 65 kDa in many tissues, especially brain.

Images

Western blot analysis of control and alkaline phosphatase-treated (AP) neonatal rat brain lysate (20 µg/lane). Blots were probed with anti-N-WASP (Lanes 1 & 2), anti-phospho-N-WASP (S484/S485) (Lanes 3 & 4), or anti-unphosphorylated-N-WASP (S484/S485) (Lanes 5 & 6).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.