

Anti-Notch 3 Homolog Antibody

Catalog # AN2188

Product Information

Application	WB, IHC
Primary Accession	<u>Q9UM47</u>
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	243631

Additional Information

Gene ID Other Names	4854 Neurogenic locus notch homolog protein 3, hN3,
Dilution	WB~~1:1000 IHC~~1:100~500
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Anti-Notch 3 Homolog Antibody is for research use only and not for use in diagnostic or therapeutic procedures.
Shipping	Blue Ice

Background

Notch signaling plays a key role in the normal development of many tissues and cell types, through diverse effects on differentiation, survival, and/or proliferation that are highly dependent on signal strength and cellular context. Members of the Notch gene family encode transmembrane receptors that are critical for various cell fate decisions. Notch family members share structural characteristics including an extracellular domain consisting of multiple epidermal growth factor-like (EGF) repeats, and an intracellular domain consisting of multiple, different domain types. Multiple human notch proteins (NOTCH1, NOTCH2, NOTCH3 and NOTCH4) have been identified and they function as a receptors for membrane bound ligands. Notch signaling is also linked to tumorigenesis as first demonstrated by the identification of a recurrent t(7;9)(q34;q34.3) chromosomal translocation involving the human Notch1 gene that is found in a small subset of human pre-T-cell acute lymphoblastic leukemias (T-ALL). Since this discovery, aberrant Notch signaling has been suggested to be involved in a wide variety of human neoplasms. Mutations in NOTCH3 have been identified as the underlying cause of cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy (CADASIL).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.