

HPRT Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1046a

Product Information

Application	WB, E
Primary Accession	P00492
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	1F8D11
Isotype	IgG2b
Calculated MW	24579
Description	The HPRT1 gene provides instructions for making an enzyme called hypoxanthine phosphoribosyltransferase 1. This enzyme allows cells to recycle purines, some of the building blocks of DNA and its chemical cousin RNA. The enzyme hypoxanthine-guanine phosphoribosyltransferase (E.C.2.4.2.8., HPRT) plays a crucial role in uric acid synthesis and purine metabolism. This enzyme catalyzes the conversion of hypoxanthine and guanine to inosine monophosphate (IMP) and guanosine monophosphate (GMP), respectively, and uses phosphoribosylpyrophosphate (PRPP) as a cosubstrate and as a source of energy. This pathway is also known as the purine salvage pathway because it allows cells to reuse purine compounds to build DNA and RNA.
Immunogen	Purified recombinant fragment of HPRT expressed in E. Coli.
Formulation	Ascitic fluid containing 0.03% sodium azide.

Additional Information

Gene ID	3251
Other Names	Hypoxanthine-guanine phosphoribosyltransferase, HGPRT, HGPRTase, 2.4.2.8, HPRT1, HPRT
Dilution	WB~~1/500 - 1/2000 E~~N/A
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	HPRT Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	HPRT1
Synonyms	HPRT
Function	Converts guanine to guanosine monophosphate, and hypoxanthine to inosine monophosphate. Transfers the 5-phosphoribosyl group from 5-phosphoribosylpyrophosphate onto the purine. Plays a central role in the generation of purine nucleotides through the purine salvage pathway.
Cellular Location	Cytoplasm.

References

1. Manjanatha MG, et.al Mutat Res. 2004 Mar 22;547(1-2):5-18.

Images




Figure 1: Western blot analysis using HPTR mouse mAb against truncated HPRT recombinant protein.




Figure 2: Immunohistochemical analysis of paraffin-embedded human normal cardiac muscle tissue, showing cytoplasmic localization using cTnI mouse mAb with DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.