

TGF beta1 Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1059a

Product Information

Application	WB, E
Primary Accession	P01137
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	4F9C10; 1B11B7
Isotype	IgG2a
Calculated MW	44325
Description	TGF beta(transforming growth factor beta), with 390-amino acid protein (about 43 kDa), is a multifunctional peptide that controls proliferation, differentiation, and other functions in many cell types. In mammals, three isoforms of TGFbeta, that is, beta1, beta2,and beta3,are known. TGF beta is one of numerous inhibitory factors produced by cancer cells that regulate antitumor immunity. TGF beta1 takes part in the local response in the course of primary lung cancer and TGFbeta1 is thought to be implicated in breast cancer progression.TGFbeta1 also plays a critical role in the downregulation of microglial responses minimizing brain inflammation and thus avoiding exacerbation of brain damage.
Immunogen	Purified recombinant fragment of TGF beta1 expressed in E. Coli.
Formulation	Ascitic fluid containing 0.03% sodium azide.

Additional Information

Gene ID	7040
Other Names	Transforming growth factor beta-1, TGF-beta-1, Latency-associated peptide, LAP, TGFB1, TGFB
Dilution	WB~~1/500 - 1/2000 E~~N/A
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TGF beta1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TGFB1 (HGNC:11766)
Synonyms	TGFB
Function	Transforming growth factor beta-1 proprotein: Precursor of the Latency-associated peptide (LAP) and Transforming growth factor beta-1 (TGF-beta-1) chains, which constitute the regulatory and active subunit of TGF-beta-1, respectively.
Cellular Location	[Latency-associated peptide]: Secreted, extracellular space, extracellular matrix
Tissue Location	Highly expressed in bone (PubMed:11746498, PubMed:17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage (PubMed:17827158)

References

1. Delphine Boche, Colm Cunningham, Fabian Docagne. Neurobiol Dis. 2006 Feb 27 2. Shizuya Saika. Lab Invest. 2006 Feb;86(2):106-15

Images




Figure 1: Western blot analysis using TGF beta1 mouse mAb against truncated TGF beta1 recombinant protein.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.