

# IKBKB Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1088a

## Product Information

---

<b>Application</b>	WB, IHC, E
<b>Primary Accession</b>	<a href="#">O14920</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	10A2C5B3
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	86564
<b>Description</b>	IKBKB(Inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase beta, also called IKK2/IKKB), is a member of the IKK complex which is composed of IKK-alpha, IKK-beta, IKK-gamma and IKAP. Phosphorylation of I-Kappa-B on a serine residue by the IKK complex frees NF-kB from I-Kappa-B and marks it for degradation via ubiquitination. IKK-beta has been shown to activate NF-kB and phosphorylate IKB-alpha and beta. Phosphorylation of 2 sites at the activation loop of IKK-beta is essential for activation of IKK by TNF and IL1. Once activated, IKK-beta autophosphorylates which in turn decreases IKK activity and prevents prolonged activation of the inflammatory response. Additionally, IKK-beta activity can also be regulated by MEKK-1.
<b>Immunogen</b>	Purified recombinant fragment of IKBKB expressed in E. Coli.
<b>Formulation</b>	Ascitic fluid containing 0.03% sodium azide.

## Additional Information

---

<b>Gene ID</b>	3551
<b>Other Names</b>	Inhibitor of nuclear factor kappa-B kinase subunit beta, I-kappa-B-kinase beta, IKK-B, IKK-beta, IkbKB, 2.7.11.10, I-kappa-B kinase 2, IKK2, Nuclear factor NF-kappa-B inhibitor kinase beta, NFKBIKB, IKBKB, IKKB
<b>Dilution</b>	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 E~~N/A
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	IKBKB Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	IKBKB
<b>Synonyms</b>	IKKB
<b>Function</b>	<p>Serine kinase that plays an essential role in the NF-kappa-B signaling pathway which is activated by multiple stimuli such as inflammatory cytokines, bacterial or viral products, DNA damages or other cellular stresses (PubMed:<a href="#">20434986</a>, PubMed:<a href="#">20797629</a>, PubMed:<a href="#">21138416</a>, PubMed:<a href="#">30337470</a>, PubMed:<a href="#">9346484</a>). Acts as a part of the canonical IKK complex in the conventional pathway of NF-kappa-B activation (PubMed:<a href="#">9346484</a>). Phosphorylates inhibitors of NF-kappa-B on 2 critical serine residues (PubMed:<a href="#">20434986</a>, PubMed:<a href="#">20797629</a>, PubMed:<a href="#">21138416</a>, PubMed:<a href="#">9346484</a>). These modifications allow polyubiquitination of the inhibitors and subsequent degradation by the proteasome (PubMed:<a href="#">20434986</a>, PubMed:<a href="#">20797629</a>, PubMed:<a href="#">21138416</a>, PubMed:<a href="#">9346484</a>). In turn, free NF-kappa-B is translocated into the nucleus and activates the transcription of hundreds of genes involved in immune response, growth control, or protection against apoptosis (PubMed:<a href="#">20434986</a>, PubMed:<a href="#">20797629</a>, PubMed:<a href="#">21138416</a>, PubMed:<a href="#">9346484</a>). In addition to the NF-kappa-B inhibitors, phosphorylates several other components of the signaling pathway including NEMO/IKBKG, NF-kappa-B subunits RELA and NFkB1, as well as IKK-related kinases TBK1 and IKBKE (PubMed:<a href="#">11297557</a>, PubMed:<a href="#">14673179</a>, PubMed:<a href="#">20410276</a>, PubMed:<a href="#">21138416</a>). IKK-related kinase phosphorylations may prevent the overproduction of inflammatory mediators since they exert a negative regulation on canonical IKKs (PubMed:<a href="#">11297557</a>, PubMed:<a href="#">20410276</a>, PubMed:<a href="#">21138416</a>). Phosphorylates FOXO3, mediating the TNF-dependent inactivation of this pro-apoptotic transcription factor (PubMed:<a href="#">15084260</a>). Also phosphorylates other substrates including NAA10, NCOA3, BCL10 and IRS1 (PubMed:<a href="#">17213322</a>, PubMed:<a href="#">19716809</a>). Phosphorylates RIPK1 at 'Ser-25' which represses its kinase activity and consequently prevents TNF- mediated RIPK1-dependent cell death (By similarity). Phosphorylates the C-terminus of IRF5, stimulating IRF5 homodimerization and translocation into the nucleus (PubMed:<a href="#">25326418</a>). Following bacterial lipopolysaccharide (LPS)-induced TLR4 endocytosis, phosphorylates STAT1 at 'Thr-749' which restricts interferon signaling and anti-inflammatory responses and promotes innate inflammatory responses (PubMed:<a href="#">38621137</a>). IKBKB-mediated phosphorylation of STAT1 at 'Thr-749' promotes binding of STAT1 to the ARID5A promoter, resulting in transcriptional activation of ARID5A and subsequent ARID5A-mediated stabilization of IL6 (PubMed:<a href="#">32209697</a>). It also promotes binding of STAT1 to the IL12B promoter and activation of IL12B transcription (PubMed:<a href="#">32209697</a>).</p>
<b>Cellular Location</b>	Cytoplasm. Nucleus. Membrane raft. Note=Colocalized with DPP4 in membrane rafts.
<b>Tissue Location</b>	Highly expressed in heart, placenta, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testis and peripheral blood

## References

1. Azoitei N,et al. Biochemistry. 2005.14;44(23): 8326-36. 2. Kumar KA,et al. Neurosci Lett. 2003.10;340(2): 139-42. 3. Peet GW,et al. J Biol Chem. 1999 Nov 12;274(46): 32655-61.

## Images




Figure 1: Western blot analysis using IKBKB mouse mAb against truncated IKBKB recombinant protein (1).




Figure 2: Immunohistochemical analysis of paraffin-embedded human colon carcinoma(A), breast carcinoma(B), kidney cell carcinoma(C), bladder carcinoma tumor(D), showing membrane and cytoplasmic localization using IKBKB mouse mAb with DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.