

# Vimentin Antibody

Purified Mouse Monoclonal Antibody  
Catalog # AO1092a

## Product Information

---

<b>Application</b>	WB, IHC, E
<b>Primary Accession</b>	<a href="#">P08670</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	9E7E7; 5G3F10; 5G3F10E5
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	53652
<b>Description</b>	<p>Vimentin is the major subunit protein of the intermediate filaments of mesenchymal cells. It is believed to be involved with the intracellular transport of proteins between the nucleus and plasma membrane. Vimentin has been implicated to be involved in the rate of steroid synthesis via its role as a storage network for steroidogenic cholesterol containing lipid droplets. Vimentin phosphorylation by a protein kinase causes the breakdown of intermediate filaments and activation of an ATP and myosin light chain dependent contractile event. This results in cytoskeletal changes that facilitate the interaction of the lipid droplets within mitochondria, and subsequent transport of cholesterol to the organelles leading to an increase in steroid synthesis. Immunohistochemical staining for Vimentin is characteristic of sarcomas (of neural, muscle and fibroblast origin) compared to carcinomas which are generally negative. Melanomas, lymphomas and vascular tumors may all stain for Vimentin. Vimentin antibodies are thus of value in the differential diagnosis of undifferentiated neoplasms and malignant tumors. They are generally used with a panel of other antibodies including those recognising cytokeratins, lymphoid markers, S100, desmin and neurofilaments.</p>
<b>Immunogen</b>	Purified recombinant fragment of Vimentin expressed in E. Coli.
<b>Formulation</b>	Ascitic fluid containing 0.03% sodium azide.

## Additional Information

---

<b>Gene ID</b>	7431
<b>Other Names</b>	Vimentin, VIM
<b>Dilution</b>	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 E~~N/A
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

## Precautions

Vimentin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

### Name

VIM ( [HGNC:12692](#) )

### Function

Vimentins are class-III intermediate filaments found in various non-epithelial cells, especially mesenchymal cells. Vimentin is attached to the nucleus, endoplasmic reticulum, and mitochondria, either laterally or terminally. Plays a role in cell directional movement, orientation, cell sheet organization and Golgi complex polarization at the cell migration front (By similarity). Protects SCRIB from proteasomal degradation and facilitates its localization to intermediate filaments in a cell contact-mediated manner (By similarity). May promote axon outgrowth and motor fiber repair via DSP-mediated recruitment to outgrowth tips (By similarity).

### Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Nucleus matrix  
{ECO:0000250|UniProtKB:P31000}. Cell membrane  
{ECO:0000250|UniProtKB:P20152}. Cell projection, axon  
{ECO:0000250|UniProtKB:P20152}

### Tissue Location

Highly expressed in fibroblasts, some expression in T- and B-lymphocytes, and little or no expression in Burkitt's lymphoma cell lines. Expressed in many hormone-independent mammary carcinoma cell lines.

## References

1. Seshadri, R., et al. *Intl. J. Cancer* 67: 353-356(1996)
2. Essa, T.M., et al. *J. Egyptian Soc. Parasitol.* 26:433-442(1996)
3. Chu, Y.W., et al. *Amer.J. Pathol.* 148: 63-69(1996)

## Images




Figure 1: Western blot analysis using Vimentin mouse mAb against truncated Vimentin recombinant protein.




Figure 2: Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue, showing cytoplasmic localization using Vimentin mouse mAb with DAB staining.




Figure 2: Immunofluorescence analysis of methanol-fixed L-02 (left) and Cos7 (right) cells using ApoM mouse mAb showing cytoplasmic and membrane localization.




Figure 2: Immunofluorescence analysis of methanol-fixed L-02(left) and COS-7(right) cells using anti-ApoM monoclonal antibody showing cytoplasmic and membrane localization.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.