10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# ALCAM Antibody

Purified Mouse Monoclonal Antibody
Catalog # AO1190a

## Product Information

| | |
|---|---|
| **Application** | IHC, E |
| **Primary Accession** | Q13740 |
| **Reactivity** | Human |
| **Host** | Mouse |
| **Clonality** | Monoclonal |
| **Clone Names** | 10F1G12 |
| **Isotype** | IgG1 |
| **Calculated MW** | 65102 |
| **Description** | ALCAM(CD166): Activated leukocyte cell adhesion molecule.CD166 is a member of the Ig superfamily and is expressed on activated T-cells, B cells and other cells including thymic epithelial cells, fibroblasts, keratinocytes and neurons. CD6 has been identified as a receptor for CD166. |
| **Immunogen** | Purified recombinant fragment of ALCAM (aa405-524) expressed in E. Coli. |
| **Formulation** | Ascitic fluid containing 0.03% sodium azide. |

## Additional Information

| | |
|---|---|
| **Gene ID** | 214 |
| **Other Names** | CD166 antigen, Activated leukocyte cell adhesion molecule, CD166, ALCAM, MEMD |
| **Dilution** | IHC~~1/200 - 1/1000 E~~N/A |
| **Storage** | Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| **Precautions** | ALCAM Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | ALCAM |
| **Synonyms** | MEMD {ECO:0000303\|PubMed:9502422} |
| **Function** | Cell adhesion molecule that mediates both heterotypic cell- cell contacts via |


| | |
|---|---|
| | its interaction with CD6, as well as homotypic cell- cell contacts (PubMed:15048703, PubMed:15496415, PubMed:16352806, PubMed:23169771, PubMed:24945728, PubMed:7760007). Promotes T-cell activation and proliferation via its interactions with CD6 (PubMed:15048703, PubMed:16352806, PubMed:24945728). Contributes to the formation and maturation of the immunological synapse via its interactions with CD6 (PubMed:15294938, PubMed:16352806). Mediates homotypic interactions with cells that express ALCAM (PubMed:15496415, PubMed:16352806). Acts as a ligand for the LILRB4 receptor, enhancing LILRB4-mediated inhibition of T cell proliferation (PubMed:29263213). Required for normal hematopoietic stem cell engraftment in the bone marrow (PubMed:24740813). Mediates attachment of dendritic cells onto endothelial cells via homotypic interaction (PubMed:23169771). Inhibits endothelial cell migration and promotes endothelial tube formation via homotypic interactions (PubMed:15496415, PubMed:23169771). Required for normal organization of the lymph vessel network. Required for normal hematopoietic stem cell engraftment in the bone marrow. Plays a role in hematopoiesis; required for normal numbers of hematopoietic stem cells in bone marrow. Promotes in vitro osteoblast proliferation and differentiation (By similarity). Promotes neurite extension, axon growth and axon guidance; axons grow preferentially on surfaces that contain ALCAM. Mediates outgrowth and pathfinding for retinal ganglion cell axons (By similarity). |
| **Cellular Location** | Cell membrane; Single-pass type I membrane protein. Cell projection, axon {ECO:0000250\|UniProtKB:Q61490}. Cell projection, dendrite {ECO:0000250\|UniProtKB:Q61490}. Note=Detected at the immunological synapse, i.e, at the contact zone between antigen-presenting dendritic cells and T-cells (PubMed:15294938, PubMed:16352806). Colocalizes with CD6 and the TCR/CD3 complex at the immunological synapse (PubMed:15294938). |
| **Tissue Location** | Detected on hematopoietic stem cells derived from umbilical cord blood (PubMed:24740813). Detected on lymph vessel endothelial cells, skin and tonsil (PubMed:23169771). Detected on peripheral blood monocytes (PubMed:15048703). Detected on monocyte- derived dendritic cells (at protein level) (PubMed:16352806). Detected at low levels in spleen, placenta, liver (PubMed:9502422). Expressed by activated T-cells, B-cells, monocytes and thymic epithelial cells (PubMed:7760007). Isoform 1 and isoform 3 are detected in vein and artery endothelial cells, astrocytes, keratinocytes and artery smooth muscle cells (PubMed:15496415). Expressed by neurons in the brain Restricted expression in tumor cell lines. Detected in highly metastasizing melanoma cell lines (PubMed:9502422) |

## References

1. Prostate. 2003 Jan 1;54(1):34-43. 2. J Clin Endocrinol Metab. 2003 Jul;88(7):3437-43. 3. J Cell Sci. 2004 Jun 1;117(Pt 13):2841-52. 4. Med Sci Monit. 2006 Aug;12(8):BR263-73.

## Images

Figure 1: Immunohistochemical analysis of paraffin-embedded human ovary carcinoma (A), kidney carcinoma (B), lung carcinoma (C) and breast carcinoma (D), showing cytoplasmic and membrane localization with DAB staining using ALCAM mouse mAb.








Figure 1: Western blot analysis using FRK mouse mAb against HEK293T cells transfected with the pCMV6-ENTRY control (1) and pCMV6-ENTRY FRK cDNA (2).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.