10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ERBB3 Antibody

Purified Mouse Monoclonal Antibody
Catalog # AO1206a

Product Information

Application	WB, IHC, E
Primary Accession	P21860
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	3F10F6
Isotype	IgG1
Calculated MW	148098
Description	ERBB3: v-erb-b2 erythroblastic leukemia viral oncogene homolog 3 (avian). This gene encodes a member of the epidermal growth factor receptor (EGFR) family of receptor tyrosine kinases. This membrane-bound protein has a neuregulin binding domain but not an active kinase domain. It therefore can bind this ligand but not convey the signal into the cell through protein phosphorylation. However, it does form heterodimers with other EGF receptor family members which do have kinase activity. Heterodimerization leads to the activation of pathways which lead to cell proliferation or differentiation. Amplification of this gene and/or overexpression of its protein have been reported in numerous cancers, including prostate, bladder, and breast tumors. Alternate transcriptional splice variants encoding different isoforms have been characterized. One isoform lacks the intermembrane region and is secreted outside the cell. This form acts to modulate the activity of the membrane-bound form. Additional splice variants have also been reported, but they have not been thoroughly characterized.
Immunogen	Purified recombinant fragment of ERBB3 (aa1175-1275) expressed in E. Coli.
Formulation	Ascitic fluid containing 0.03% sodium azide.

Additional Information

Gene ID	2065
Other Names	Receptor tyrosine-protein kinase erbB-3, 2.7.10.1, Proto-oncogene-like protein c-ErbB-3, Tyrosine kinase-type cell surface receptor HER3, ERBB3, HER3
Dilution	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 E~~N/A
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ERBB3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	ERBB3
Synonyms	HER3
Function	Tyrosine kinase receptor that, despite substitutions of key catalytic residues in the kinase domain, retains residual catalytic activity enabling autophosphorylation (PubMed:9148746, PubMed:20351256). Functions primarily as a heterodimer, preferentially partnering with ERBB2/HER2, which phosphorylates ERBB3 on specific tyrosine residues and provides the kinase activity required for downstream signaling (PubMed:22983903). Binds neuregulins NRG1 and NRG2 through its extracellular domain, which induces conformational change from a tethered autoinhibited state to an extended conformation competent for dimerization (PubMed:20682778). Upon heterodimerization and trans-phosphorylation, acts as the most efficient activator of the PI3K/AKT pathway within the ErbB family (PubMed:22983903). Also activates the RAS/MAPK/ERK cascade to promote cell proliferation (PubMed:22983903). Plays an essential role in cardiac development; required for normal formation of the endocardial cushion and ventricular trabeculation during embryogenesis (Potential). Critical for peripheral nervous system development, where neuregulin-1/ERBB3 signaling drives Schwann cell survival, proliferation, and differentiation, and is required for axonal myelination (By similarity). Involved in epithelial morphogenesis and mammary gland development (By similarity). Also involved in the regulation of myeloid cell differentiation (PubMed:27416908). Participates in a regulatory feedback loop whereby pathway activation induces transcriptional up-regulation of ERBB3 expression (PubMed:21368164). Participates in vesicular sorting by stabilizing the ARF6-GGA3-RABEP1 endosomal sorting complex, to promote early recycling of TFRC and ITGB1 (PubMed:41348103). May also be activated by additional ligands including CSPG5 and EGFL6; the latter triggers assembly of an ERBB2/ERBB3/ITGB3 signaling complex leading to ERK phosphorylation and cell proliferation (PubMed:15358134, PubMed:41151154).
Cellular Location	[Isoform 1]: Cell membrane; Single-pass type I membrane protein
Tissue Location	Epithelial tissues and brain.

References

1. Cancer Sci. 2007 Sep;98(9):1498-503. 2. Breast Cancer Res. 2008;10(1):R2.

Images

Figure 1: Western blot analysis using ERBB3 mouse mAb against truncated Trx-ERBB3 recombinant protein (1) and MBP-ERBB3 (aa1175-1275) recombinant protein (2).






Figure 2: Immunohistochemical analysis of paraffin-embedded human cerebra (left) and breast carcinoma tissue (right),showing cytoplasmic and membrane location with DAB staining using ERBB3 mouse mAb.




Figure 2: Confocal immunofluorescence analysis of Hela cells using HDAC3 mouse mAb (green). Red: Actin filaments have been labeled with DY-554 phalloidin.




Figure 2: Confocal immunofluorescence analysis of Hela cells using anti-HDAC3 mAb (green). Red: Actin filaments have been labeled with DY-554 phalloidin.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.