

# PGR Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1219a

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">P06401</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	8A11H1
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	98981
<b>Description</b>	PGR: progesterone receptor. This gene encodes a member of the steroid receptor superfamily. The encoded protein mediates the physiological effects of progesterone, which plays a central role in reproductive events associated with the establishment and maintenance of pregnancy. This gene uses two distinct promoters and translation start sites in the first exon to produce two isoforms, A and B. The two isoforms are identical except for the additional 165 amino acids found in the N-terminus of isoform A only, and mediate their own response genes and physiologic effects with little overlap. The location of transcription initiation for isoform B has not been clearly determined.
<b>Immunogen</b>	Purified recombinant fragment of PGR (aa730-871) expressed in E. Coli.
<b>Formulation</b>	Ascitic fluid containing 0.03% sodium azide.

## Additional Information

---

<b>Gene ID</b>	5241
<b>Other Names</b>	Progesterone receptor, PR, Nuclear receptor subfamily 3 group C member 3, PGR, NR3C3
<b>Dilution</b>	WB~~1/500 - 1/2000 E~~N/A
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	PGR Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	PGR
<b>Synonyms</b>	NR3C3
<b>Function</b>	The steroid hormones and their receptors are involved in the regulation of eukaryotic gene expression and affect cellular proliferation and differentiation in target tissues. Depending on the isoform, progesterone receptor functions as a transcriptional activator or repressor.
<b>Cellular Location</b>	Nucleus. Cytoplasm. Note=Nucleoplasmic shuttling is both hormone- and cell cycle-dependent. On hormone stimulation, retained in the cytoplasm in the G(1) and G(2)/M phases [Isoform 4]; Mitochondrion outer membrane
<b>Tissue Location</b>	In reproductive tissues the expression of isoform A and isoform B varies as a consequence of developmental and hormonal status. Isoform A and isoform B are expressed in comparable levels in uterine glandular epithelium during the proliferative phase of the menstrual cycle. Expression of isoform B but not of isoform A persists in the glands during mid-secretory phase. In the stroma, isoform A is the predominant form throughout the cycle. Heterogeneous isoform expression between the glands of the endometrium basalis and functionalis is implying region-specific responses to hormonal stimuli

## References

1. Cancer Sci. 2006 Dec;97(12):1308. 2. Mol Endocrinol. 2006 Nov;20(11):2656-70. 3. Mol Endocrinol. 2007 Jan;21(1):106-25. 4. J Clin Endocrinol Metab. 2007 Nov;92(11):4459-66.

## Images




Figure 1: Western blot analysis using PGR mouse mAb against truncated MBP-PGR recombinant protein (1) and truncated Trx-PGR(aa730-871) recombinant protein (2).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.