

# Pirh2 Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1358a

## Product Information

---

<b>Application</b>	WB, IHC, FC, ICC, E
<b>Primary Accession</b>	<a href="#">Q96PM5</a>
<b>Reactivity</b>	Human, Rat
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	1H10
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	30110
<b>Description</b>	Pirh 2 (P53 induced RING-H2 protein), also known as RCHY1, it forms dimers through its N- and C-terminus in cells. The Pirh2 has ubiquitin-protein ligase activity and it binds with p53 and promotes the ubiquitin-mediated proteosomal degradation of p53. The Pirh2 is oncogenic because loss of p53 function contributes directly to malignant tumor development. Pirh2 expression decreases the level of p53, and a decrease of endogenous Pirh2 expression increases p53 levels. Pirh2 is therefore considered, together with MDM2, to act as a negative regulator of p53 function.
<b>Immunogen</b>	Purified recombinant fragment of human Pirh2 expressed in E. Coli.
<b>Formulation</b>	Ascitic fluid containing 0.03% sodium azide.

## Additional Information

---

<b>Gene ID</b>	25898
<b>Other Names</b>	RING finger and CHY zinc finger domain-containing protein 1, 6.3.2.-, Androgen receptor N-terminal-interacting protein, CH-rich-interacting match with PLAG1, E3 ubiquitin-protein ligase Pirh2, RING finger protein 199, Zinc finger protein 363, p53-induced RING-H2 protein, hPirh2, RCHY1, ARNIP, CHIMP, PIRH2, RNF199, ZNF363
<b>Dilution</b>	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 FC~~1/200 - 1/400 ICC~~N/A E~~N/A
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Pirh2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

Name	RCHY1
Function	E3 ubiquitin-protein ligase that mediates ubiquitination of target proteins, including p53/TP53, TP73, HDAC1 and CDKN1B (PubMed: <a href="#">16914734</a> , PubMed: <a href="#">17721809</a> , PubMed: <a href="#">18006823</a> , PubMed: <a href="#">19043414</a> , PubMed: <a href="#">19483087</a> , PubMed: <a href="#">21994467</a> ). Mediates ubiquitination and degradation of p53/TP53; preferentially acts on tetrameric p53/TP53 (PubMed: <a href="#">19043414</a> , PubMed: <a href="#">19483087</a> ). Catalyzes monoubiquitinates the translesion DNA polymerase POLH (PubMed: <a href="#">21791603</a> ). Involved in the ribosome-associated quality control (RQC) pathway, which mediates the extraction of incompletely synthesized nascent chains from stalled ribosomes: RCHY1 acts downstream of NEMF and recognizes CAT tails associated with stalled nascent chains, leading to their ubiquitination and degradation (PubMed: <a href="#">33909987</a> ).
Cellular Location	Nucleus. Nucleus speckle. Cytoplasm

## References

1. Biochem Biophys Res Commun. 2007 Dec 14;364(2):344-50. 2. J Natl Cancer Inst. 2004 Nov 17;96(22):1718-21. 3. Exp Cell Res. 2006 Oct 15;312(17):3370-8. 4. Cell. 2003 Mar 21;112(6):779-91.

## Images




Figure 1: Western blot analysis using Pirh2 mouse mAb against Hela (1), A549 (2), MCF-7 (3) and PC-12 (4) cell lysate.




Figure 2: Immunohistochemical analysis of paraffin-embedded human Tonsil tissues using anti-Pirh2 mouse mAb

Figure 3: Flow cytometric analysis of PC-12 cells using anti-Pirh2 mAb (blue) and negative control (red).





Figure 4: Immunofluorescence analysis of Hela cells using Pirh2 mouse mAb (green). Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.