

PAX8 Antibody

Purified Mouse Monoclonal Antibody Catalog # AO1470a

Product Information

Application WB, E
Primary Accession Q71U36
Reactivity Human
Host Mouse
Clonality Monoclonal

Clone Names 2D4 Isotype IgG1 Calculated MW 50136

Description This gene encodes a member of the paired box (PAX) family of transcription

factors. Members of this gene family typically encode proteins that contain a paired box domain, an octapeptide, and a paired-type homeodomain. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas and atypical follicular thyroid adenomas. Tissue specificity: Expressed in the excretory system, thyroid gland and Wilms tumors. ABCAM: This protein is a member of the paired box (PAX) family of transcription factors, typically containing a paired box domain, an octapeptide, and a paired-type homeodomain. This family plays critical roles during fetal development and cancer growth. The specific function of the PAX8 is unknown but it may involve kidney cell differentiation, thyroid development, or thyroid dysgenesis. Alternative splicing in the gene by inclusion or exclusion of exons 7 and/or 8 has produced several known products but the biological significance of the variants is unknown. Several other splice variants have been proposed but the

full nature of these products has not been described. Pax8 is also a marker of

otic progenitor cells.

Immunogen Purified recombinant fragment of human PAX8 expressed in E. Coli.

Formulation Ascitic fluid containing 0.03% sodium azide.

Additional Information

Gene ID 7846

Other Names Tubulin alpha-1A chain, Alpha-tubulin 3, Tubulin B-alpha-1, Tubulin alpha-3

chain, TUBA1A, TUBA3

Dilution WB~~1/500 - 1/2000 E~~N/A

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

therapeutic procedures.

Protein Information

Name TUBA1A

Synonyms TUBA3

Function Tubulin is the major constituent of microtubules, a cylinder consisting of

laterally associated linear protofilaments composed of alpha- and

beta-tubulin heterodimers. Microtubules grow by the addition of GTP-tubulin dimers to the microtubule end, where a stabilizing cap forms. Below the cap,

tubulin dimers are in GDP-bound state, owing to GTPase activity of

alpha-tubulin.

Cellular Location Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, flagellum axoneme

{ECO:0000250 | UniProtKB:P68369}

Tissue Location Expressed at a high level in fetal brain.

References

1. Thyroid. 2009 Jan;19(1):61-8. 2. Cancer Genet Cytogenet. 2010 Jan 1;196(1):7-13. 3. Cancer Cytopathol. 2010 Oct 25;118(5):298-302.

Images




Figure 1: Western blot analysis using PAX8 mouse mAb against Hela (1),HEK293 (2) and Raji (3) cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.