

FOXD3 Antibody

Purified Mouse Monoclonal Antibody Catalog # AO1480a

Product Information

Application WB, E Primary Accession Q9UJU5

Reactivity Human, Mouse, Monkey

Host Mouse **Clonality** Monoclonal

Clone Names5G9IsotypeIgG1Calculated MW47630

Description FoxD3 is a member of the Forkhead Box family and is characterized by a

winged-helix DNA-binding structure and the important role it plays in embryonic development . This transcriptional regulator is required for the maintenance of pluripotency in the pre-implantation and peri-implantation stages of mouse embryonic development and is also required for trophoblast formation . FoxD3 is required for the maintenance of the mammalian neural crest; FoxD3(-/-) mouse embryos fail around the time of implantation with loss of neural crest-derived structures . FoxD3 also forms a regulatory network with Oct-4 and NANOG to maintain the pluripotency of ES cells .Promotes development of neural crest cells from neural tube progenitors. Restricts neural progenitor cells to the neural crest lineage while suppressing interneuron differentiation. Required for maintenance of pluripotent cells in the pre-implantation and peri-implantation stages of embryogenesis .Tissue specificity: Expressed in chronic myeloid leukemia, Jurkat T-cell leukemia and teratocarcinoma cell lines, but not in any other cell lines or normal tissues

examined.

Immunogen Purified recombinant fragment of human FOXD3 expressed in E. Coli.

Formulation Ascitic fluid containing 0.03% sodium azide.

Additional Information

Gene ID 27022

Other Names Forkhead box protein D3, HNF3/FH transcription factor genesis, FOXD3, HFH2

Dilution WB~~1/500 - 1/2000 E~~N/A

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions FOXD3 Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

Protein Information

Name FOXD3

Synonyms HFH2

Function Binds to the consensus sequence 5'-A[AT]T[AG]TTTGTTT-3' and acts as a

transcriptional repressor (PubMed:<u>11891324</u>). Also acts as a transcriptional activator (PubMed:<u>11891324</u>). Negatively regulates transcription of transcriptional repressor RHIT/ZNF205 (PubMed:<u>22306510</u>). Promotes

development of neural crest cells from neural tube progenitors

(PubMed:<u>11891324</u>). Restricts neural progenitor cells to the neural crest lineage while suppressing interneuron differentiation (PubMed:<u>11891324</u>). Required for maintenance of pluripotent cells in the pre-implantation and

peri-implantation stages of embryogenesis (PubMed:11891324).

Cellular Location Nucleus.

Tissue Location Expressed in chronic myeloid leukemia, Jurkat T- cell leukemia and

teratocarcinoma cell lines, but not in any other cell lines or normal tissues

examined.

References

1. Blood. 1993 Jun 1;81(11):2854-9. 2. Am J Hum Genet. 2001 Mar;68(3):627-41.

Images




Figure 1: Western blot analysis using FOXD3 mouse mAb against NTERA-1 (1), HUVE-12 (2), HEK293 (3), Hela (4), Jurkat (5), NIH/3T3 (6), K562 (7), RAW264.7 (8) and COS7 (9) cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.