

SUZ12 Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1494a

Product Information

| | |
|--------------------------|--|
| Application | WB, ICC, E |
| Primary Accession | Q15022 |
| Reactivity | Human |
| Host | Mouse |
| Clonality | Monoclonal |
| Clone Names | 2C11 |
| Isotype | IgG1 |
| Calculated MW | 83055 |
| Description | This zinc finger gene has been identified at the breakpoints of a recurrent chromosomal translocation reported in endometrial stromal sarcoma. Recombination of these breakpoints results in the fusion of this gene and JAZF1. The protein encoded by this gene contains a zinc finger domain in the C terminus of the coding region. (Provided by RefSeq) SUZ12 is overexpressed in several human tumors, including tumors of the colon, breast and liver. Tissue specificity: Overexpressed in breast and colon cancer. |
| Immunogen | Purified recombinant fragment of human SUZ12 expressed in E. Coli. |
| Formulation | Ascitic fluid containing 0.03% sodium azide. |

Additional Information

| | |
|--------------------|---|
| Gene ID | 23512 |
| Other Names | Polycomb protein SUZ12, Chromatin precipitated E2F target 9 protein, ChET 9 protein, Joined to JAZF1 protein, Suppressor of zeste 12 protein homolog, SUZ12, CHET9, JJAZ1, KIAA0160 |
| Dilution | WB~~1/500 - 1/2000 ICC~~N/A E~~N/A |
| Storage | Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | SUZ12 Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|-------------|-------|
| Name | SUZ12 |
|-------------|-------|

Synonyms

CHET9, JJAZ1, KIAA0160

Function

Polycomb group (PcG) protein. Component of the PRC2 complex, which methylates 'Lys-9' (H3K9me) and 'Lys-27' (H3K27me) of histone H3, leading to transcriptional repression of the affected target gene (PubMed:[15225548](#), PubMed:[15231737](#), PubMed:[15385962](#), PubMed:[16618801](#), PubMed:[17344414](#), PubMed:[18285464](#), PubMed:[28229514](#), PubMed:[29499137](#), PubMed:[31959557](#)). The PRC2 complex may also serve as a recruiting platform for DNA methyltransferases, thereby linking two epigenetic repression systems (PubMed:[12351676](#), PubMed:[12435631](#), PubMed:[15099518](#), PubMed:[15225548](#), PubMed:[15385962](#), PubMed:[15684044](#), PubMed:[16431907](#), PubMed:[18086877](#), PubMed:[18285464](#)). Genes repressed by the PRC2 complex include HOXC8, HOXA9, MYT1 and CDKN2A (PubMed:[15231737](#), PubMed:[16618801](#), PubMed:[17200670](#), PubMed:[31959557](#)).

Cellular Location

Nucleus Note=Localizes to chromatin as part of the PRC2 complex

Tissue Location

Overexpressed in breast and colon cancer.

References

1. Genes Dev. 2008 May 15;22(10):1345-55. 2. Proc Natl Acad Sci U S A. 2007 Dec 11;104(50):20001-6.

Images




Figure 1: Western blot analysis using SUZ12 mAb against HEK293 (1) and SUZ12(AA: 533-739)-hIgGFc transfected HEK293 (2) cell lysate.





Figure 2: Immunofluorescence analysis of U251 cells using SUZ12 mouse mAb (green). Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.

Figure 3: Immunofluorescence analysis of MCF-7 cells using SUZ12 mouse mAb (green). Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.