

# CA9 Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1500a

## Product Information

---

<b>Application</b>	WB, IHC, FC, E
<b>Primary Accession</b>	<a href="#">Q16790</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	2D3
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	49698
<b>Description</b>	CA IX is a transmembrane protein and the only tumor-associated carbonic anhydrase isoenzyme known. It is expressed in all clear-cell renal cell carcinoma, but is not detected in normal kidney or most other normal tissues. It may be involved in cell proliferation and transformation. Reversible hydration of carbon dioxide. Participates in pH regulation. May be involved in the control of cell proliferation and transformation. Appears to be a novel specific biomarker for a cervical neoplasia. Tissue specificity: Expressed primarily in carcinoma cells lines. Expression is restricted to very few normal tissues and the most abundant expression is found in the epithelial cells of gastric mucosa.
<b>Immunogen</b>	Purified recombinant fragment of human CA9 expressed in E. Coli.
<b>Formulation</b>	Ascitic fluid containing 0.03% sodium azide.

## Additional Information

---

<b>Gene ID</b>	768
<b>Other Names</b>	Carbonic anhydrase 9, 4.2.1.1, Carbonate dehydratase IX, Carbonic anhydrase IX, CA-IX, CAIX, Membrane antigen MN, P54/58N, Renal cell carcinoma-associated antigen G250, RCC-associated antigen G250, pMW1, CA9, G250, MN
<b>Dilution</b>	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 FC~~1/200 - 1/400 E~~N/A
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	CA9 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

<b>Name</b>	CA9
<b>Synonyms</b>	G250, MN
<b>Function</b>	Catalyzes the interconversion between carbon dioxide and water and the dissociated ions of carbonic acid (i.e. bicarbonate and hydrogen ions).
<b>Cellular Location</b>	Nucleus. Nucleus, nucleolus. Cell membrane; Single-pass type I membrane protein. Cell projection, microvillus membrane; Single-pass type I membrane protein. Note=Found on the surface microvilli and in the nucleus, particularly in nucleolus
<b>Tissue Location</b>	Expressed primarily in carcinoma cells lines. Expression is restricted to very few normal tissues and the most abundant expression is found in the epithelial cells of gastric mucosa

## References

1. Br J Cancer. 2008 Sep 2;99(5):727-33. 2. Pathol Res Pract. 2009;205(1):1-9.

## Images




Figure 1: Western blot analysis using CA9 mouse mAb against Hela (1) and A549 (2) cell lysate.




Figure 2: Immunohistochemical analysis of paraffin-embedded stomach tissues (left) and colon tissues (right) using CA9 mouse mAb with DAB staining.




Figure 3: Flow cytometric analysis of NTERA-2 cells using CA9 mouse mAb (green) and negative control (purple).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.