

# HIF1A Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1546a

## Product Information

---

<b>Application</b>	WB, IHC, ICC, E
<b>Primary Accession</b>	<a href="#">Q16665</a>
<b>Reactivity</b>	Human, Mouse, Monkey
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	1A3
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	92670
<b>Description</b>	<p>Hypoxia-inducible factor-1 (HIF1) is a transcription factor found in mammalian cells cultured under reduced oxygen tension that plays an essential role in cellular and systemic homeostatic responses to hypoxia. HIF1 is a heterodimer composed of an alpha subunit and a beta subunit. The beta subunit has been identified as the aryl hydrocarbon receptor nuclear translocator (ARNT). This gene encodes the alpha subunit of HIF-1. Overexpression of a natural antisense transcript (aHIF) of this gene has been shown to be associated with nonpapillary renal carcinomas. Two alternative transcripts encoding different isoforms have been identified. (provided by RefSeq) Tissue specificity: Expressed in most tissues with highest levels in kidney and heart. Overexpressed in the majority of common human cancers and their metastases, due to the presence of intratumoral hypoxia and as a result of mutations in genes encoding oncoproteins and tumor suppressors.</p>
<b>Immunogen</b>	Purified recombinant fragment of human HIF1A expressed in E. Coli.
<b>Formulation</b>	Ascitic fluid containing 0.03% sodium azide.

## Additional Information

---

<b>Gene ID</b>	3091
<b>Other Names</b>	Hypoxia-inducible factor 1-alpha, HIF-1-alpha, HIF1-alpha, ARNT-interacting protein, Basic-helix-loop-helix-PAS protein MOP1, Class E basic helix-loop-helix protein 78, bHLHe78, Member of PAS protein 1, PAS domain-containing protein 8, HIF1A, BHLHE78, MOP1, PASD8
<b>Dilution</b>	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 ICC~~N/A E~~1/10000
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	HIF1A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	HIF1A {ECO:0000303   PubMed:7539918, ECO:0000312   HGNC:HGNC:4910}
<b>Function</b>	Functions as a master transcriptional regulator of the adaptive response to hypoxia (PubMed: <a href="#">11292861</a> , PubMed: <a href="#">11566883</a> , PubMed: <a href="#">15465032</a> , PubMed: <a href="#">16973622</a> , PubMed: <a href="#">17610843</a> , PubMed: <a href="#">18658046</a> , PubMed: <a href="#">20624928</a> , PubMed: <a href="#">22009797</a> , PubMed: <a href="#">30125331</a> , PubMed: <a href="#">9887100</a> ). Under hypoxic conditions, activates the transcription of over 40 genes, including erythropoietin, glucose transporters, glycolytic enzymes, vascular endothelial growth factor, HILPDA, and other genes whose protein products increase oxygen delivery or facilitate metabolic adaptation to hypoxia (PubMed: <a href="#">11292861</a> , PubMed: <a href="#">11566883</a> , PubMed: <a href="#">15465032</a> , PubMed: <a href="#">16973622</a> , PubMed: <a href="#">17610843</a> , PubMed: <a href="#">20624928</a> , PubMed: <a href="#">22009797</a> , PubMed: <a href="#">30125331</a> , PubMed: <a href="#">9887100</a> ). Plays an essential role in embryonic vascularization, tumor angiogenesis and pathophysiology of ischemic disease (PubMed: <a href="#">22009797</a> ). Heterodimerizes with ARNT; heterodimer binds to core DNA sequence 5'-TACGTG-3' within the hypoxia response element (HRE) of target gene promoters (By similarity). Activation requires recruitment of transcriptional coactivators such as CREBBP and EP300 (PubMed: <a href="#">16543236</a> , PubMed: <a href="#">9887100</a> ). Activity is enhanced by interaction with NCOA1 and/or NCOA2 (PubMed: <a href="#">10594042</a> ). Interaction with redox regulatory protein APEX1 seems to activate CTAD and potentiates activation by NCOA1 and CREBBP (PubMed: <a href="#">10202154</a> , PubMed: <a href="#">10594042</a> ). Involved in the axonal distribution and transport of mitochondria in neurons during hypoxia (PubMed: <a href="#">19528298</a> ).
<b>Cellular Location</b>	Cytoplasm. Nucleus. Nucleus speckle {ECO:0000250   UniProtKB:Q61221}. Note=Colocalizes with HIF3A in the nucleus and speckles (By similarity). Cytoplasmic in normoxia, nuclear translocation in response to hypoxia (PubMed:9822602) {ECO:0000250   UniProtKB:Q61221, ECO:0000269   PubMed:9822602}
<b>Tissue Location</b>	Expressed in most tissues with highest levels in kidney and heart. Overexpressed in the majority of common human cancers and their metastases, due to the presence of intratumoral hypoxia and as a result of mutations in genes encoding oncoproteins and tumor suppressors. A higher level expression seen in pituitary tumors as compared to the pituitary gland.

## References

---

1. Int J Radiat Oncol Biol Phys. 2008 Dec 1;72(5):1551-9. 2. Eur J Appl Physiol. 2009 Mar;105(4):515-24.

## Images

---

Figure 1: Western blot analysis using HIF1A mouse mAb against Cos7 (1), Hela (2), Jurkat (3), RAJI (4) and NIH/3T3 (5) cell lysate.




Figure 2: Immunohistochemical analysis of paraffin-embedded liver cancer tissues (left) and lung cancer tissues (right) using HIF1A mouse mAb with DAB staining.

Figure 3: Immunohistochemical analysis of paraffin-embedded stomach cancer tissues (left) and brain tumor tissues (right) using HIF1A mouse mAb with DAB staining.

Figure 4: Immunofluorescence analysis of HeLa cells using HIF1A mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.