

# ABCG2 Antibody

Purified Mouse Monoclonal Antibody  
Catalog # AO1569a

## Product Information

---

<b>Application</b>	WB, ICC, E
<b>Primary Accession</b>	<a href="#">Q9UNQ0</a>
<b>Reactivity</b>	Human, Mouse, Monkey
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	1H2
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	72314
<b>Description</b>	<p>The membrane-associated protein encoded by this gene is included in the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the White subfamily. Alternatively referred to as a breast cancer resistance protein, this protein functions as a xenobiotic transporter which may play a major role in multi-drug resistance. It likely serves as a cellular defense mechanism in response to mitoxantrone and anthracycline exposure. Significant expression of this protein has been observed in the placenta, which may suggest a potential role for this molecule in placenta tissue. Tissue specificity: Highly expressed in placenta. Low expression in small intestine, liver and colon.</p>
<b>Immunogen</b>	Purified recombinant fragment of human ABCG2 expressed in E. Coli.
<b>Formulation</b>	Ascitic fluid containing 0.03% sodium azide.

## Additional Information

---

<b>Gene ID</b>	9429
<b>Other Names</b>	ATP-binding cassette sub-family G member 2, Breast cancer resistance protein, CDw338, Mitoxantrone resistance-associated protein, Placenta-specific ATP-binding cassette transporter, Urate exporter, CD338, ABCG2, ABCP, BCRP, BCRP1, MXR
<b>Dilution</b>	WB~~1/500 - 1/2000 ICC~~N/A E~~1/10000
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	ABCG2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	ABCG2 ( <a href="#">HGNC:74</a> )
<b>Synonyms</b>	ABCP, BCRP, BCRP1, MXR
<b>Function</b>	<p>Broad substrate specificity ATP-dependent transporter of the ATP-binding cassette (ABC) family that actively extrudes a wide variety of physiological compounds, dietary toxins and xenobiotics from cells (PubMed:<a href="#">11306452</a>, PubMed:<a href="#">12958161</a>, PubMed:<a href="#">19506252</a>, PubMed:<a href="#">20705604</a>, PubMed:<a href="#">28554189</a>, PubMed:<a href="#">30405239</a>, PubMed:<a href="#">31003562</a>). Involved in porphyrin homeostasis, mediating the export of protoporphyrin IX (PPIX) from both mitochondria to cytosol and cytosol to extracellular space, it also functions in the cellular export of heme (PubMed:<a href="#">20705604</a>, PubMed:<a href="#">23189181</a>). Also mediates the efflux of sphingosine-1-P from cells (PubMed:<a href="#">20110355</a>). Acts as a urate exporter functioning in both renal and extrarenal urate excretion (PubMed:<a href="#">19506252</a>, PubMed:<a href="#">20368174</a>, PubMed:<a href="#">22132962</a>, PubMed:<a href="#">31003562</a>, PubMed:<a href="#">36749388</a>). In kidney, it also functions as a physiological exporter of the uremic toxin indoxyl sulfate (By similarity). Also involved in the excretion of steroids like estrone 3-sulfate/E1S, 3beta-sulfoxy-androst-5-en-17-one/DHEAS, and other sulfate conjugates (PubMed:<a href="#">12682043</a>, PubMed:<a href="#">28554189</a>, PubMed:<a href="#">30405239</a>). Mediates the secretion of the vitamins riboflavin and biotin into milk (By similarity). Involved in the excretion of the riboflavin-derived compound lumichrome into the intestinal lumen and in its secretion into milk (PubMed:<a href="#">39337371</a>). Extrudes pheophorbide a, a phototoxic porphyrin catabolite of chlorophyll, reducing its bioavailability (By similarity). Plays an important role in the exclusion of xenobiotics from the brain (Probable). It confers to cells a resistance to multiple drugs and other xenobiotics including mitoxantrone, pheophorbide, camptothecin, methotrexate, azidothymidine, and the anthracyclines daunorubicin and doxorubicin, through the control of their efflux (PubMed:<a href="#">11306452</a>, PubMed:<a href="#">12477054</a>, PubMed:<a href="#">15670731</a>, PubMed:<a href="#">18056989</a>, PubMed:<a href="#">31254042</a>). In placenta, it limits the penetration of drugs from the maternal plasma into the fetus (By similarity). May play a role in early stem cell self-renewal by blocking differentiation (By similarity). In inflammatory macrophages, exports itaconate from the cytosol to the extracellular compartment and limits the activation of TFEB-dependent lysosome biogenesis involved in antibacterial innate immune response.</p>
<b>Cellular Location</b>	Cell membrane; Multi-pass membrane protein. Apical cell membrane; Multi-pass membrane protein. Mitochondrion membrane; Multi-pass membrane protein. Note=Enriched in membrane lipid rafts
<b>Tissue Location</b>	Highly expressed in placenta (PubMed:9850061). Low expression in small intestine, liver and colon (PubMed:9861027) Expressed in brain (at protein level) (PubMed:12958161)

---

## References

1. Carcinogenesis. 2008 Dec;29(12):2289-97.
2. Pharm Res. 2009 Feb;26(2):449-58.

---

## Images

Figure 1: Western blot analysis using ABCG2 mouse mAb against HepG2 (1), Cos7 (2), Jurkat (3) and NIH/3T3 (4) cell





Figure 2: Immunofluorescence analysis of Hela cells using ABCG2 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.