

KID Antibody

Purified Mouse Monoclonal Antibody Catalog # AO1627a

Product Information

Application WB, IHC, FC, E
Primary Accession Q14807
Reactivity Human
Host Mouse
Clonality Monoclonal

Clone Names1E3IsotypeIgG1Calculated MW73262

Description The protein encoded by this gene is a member of kinesin-like protein family.

This family of proteins are microtubule-dependent molecular motors that transport organelles within cells and move chromosomes during cell division. The C-terminal half of this protein has been shown to bind DNA. Studies with the Xenopus homolog suggests its essential role in metaphase chromosome

alignment and maintenance.

Immunogen Purified recombinant fragment of human KID expressed in E. Coli.

Formulation Ascitic fluid containing 0.03% sodium azide.

Additional Information

Gene ID 3835

Other Names Kinesin-like protein KIF22, Kinesin-like DNA-binding protein, Kinesin-like

protein 4, KIF22, KID, KNSL4

Dilution WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 FC~~1/200 - 1/400 E~~1/10000

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions KID Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

Protein Information

Name KIF22

Synonyms KID, KNSL4

Function Kinesin family member that is involved in spindle formation and the

movements of chromosomes during mitosis and meiosis. Binds to microtubules and to DNA (By similarity). Plays a role in congression of

laterally attached chromosomes in NDC80-depleted cells (PubMed:25743205).

Cellular Location Nucleus. Cytoplasm, cytoskeleton

Tissue Location Expressed in bone, cartilage, joint capsule, ligament, skin, and primary

cultured chondrocytes

References

1. Cell. 2008 Mar 7;132(5):771-82. 2. Retrovirology. 2009 May 19;6:47.

Images




Figure 1: Western blot analysis using KID mAb against human KID (AA: 225-419) recombinant protein. (Expected MW is 47 kDa)

Figure 2: Western blot analysis using KID mAb against HEK293 (1) and KID(AA: 225-419)-hIgGFc transfected HEK293 (2) cell lysate.





Figure 3: Immunohistochemical analysis of paraffin-embedded rectum cancer tissues using KID mouse mAb with DAB staining.

Figure 4: Immunohistochemical analysis of paraffin-embedded colon cancer tissues using KID mouse mAb with DAB staining.






Figure 5: Flow cytometric analysis of NIH/3T3 cells using KID mouse mAb (blue) and negative control (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.