

# TNNI2 Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1628a

## Product Information

---

<b>Application</b>	WB, IHC, FC, E
<b>Primary Accession</b>	<a href="#">P48788</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone Names</b>	2F12A8
<b>Isotype</b>	IgG1
<b>Calculated MW</b>	21339
<b>Description</b>	This gene encodes a fast-twitch skeletal muscle protein, a member of the troponin I gene family, and a component of the troponin complex including troponin T, troponin C and troponin I subunits. The troponin complex, along with tropomyosin, is responsible for the calcium-dependent regulation of striated muscle contraction. Mouse studies show that this component is also present in vascular smooth muscle and may play a role in regulation of smooth muscle function. In addition to muscle tissues, this protein is found in corneal epithelium, cartilage where it is an inhibitor of angiogenesis to inhibit tumor growth and metastasis, and mammary gland where it functions as a co-activator of estrogen receptor-related receptor alpha. This protein also suppresses tumor growth in human ovarian carcinoma. Mutations in this gene cause myopathy and distal arthrogryposis type 2B. Alternatively spliced transcript variants have been found for this gene.
<b>Immunogen</b>	Purified recombinant fragment of human TNNI2 expressed in E. Coli.
<b>Formulation</b>	Ascitic fluid containing 0.03% sodium azide.

## Additional Information

---

<b>Gene ID</b>	7136
<b>Other Names</b>	Troponin I, fast skeletal muscle, Troponin I, fast-twitch isoform, TNNI2
<b>Dilution</b>	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 FC~~1/200 - 1/400 E~~1/10000
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	TNNI2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

Name	TNNI2
Function	Troponin I is the inhibitory subunit of troponin, the thin filament regulatory complex which confers calcium-sensitivity to striated muscle actomyosin ATPase activity.

## References

1. Am J Hum Genet. 2009 Nov;85(5):628-42. 2. Cell Motil Cytoskeleton. 2008 Aug;65(8):652-61.

## Images




Figure 1: Western blot analysis using TNNI2 mAb against human TNNI2 (AA: 1-182) recombinant protein.(Expected MW is 46.8 kDa)




Figure 2: Western blot analysis using TNNI2 mAb against HEK293 (1) and TNNI2(AA: 1-182)-hIgGfc transfected HEK293 (2) cell lysate.




Figure 3: Immunohistochemical analysis of paraffin-embedded cardiac muscle tissues using TNNI2 mouse mAb with DAB staining.




Figure 4: Immunohistochemical analysis of paraffin-embedded striated muscle tissues using TNNI2 mouse mAb with DAB staining.




Figure 5: Flow cytometric analysis of NIH/3T3 cells using TNNI2 mouse mAb (blue) and negative control (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.