

CYP3A4 Antibody

Purified Mouse Monoclonal Antibody

Catalog # AO1696a

Product Information

Application	WB, IHC, FC, ICC, E
Primary Accession	P08684
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Clone Names	3H8
Isotype	IgG1
Calculated MW	57343
Description	This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases that catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by glucocorticoids and some pharmacological agents. This enzyme is involved in the metabolism of approximately half the drugs in use today, including acetaminophen, codeine, cyclosporin A, diazepam and erythromycin. The enzyme also metabolizes some steroids and carcinogens. This gene is part of a cluster of cytochrome P450 genes on chromosome 7q21.1. Previously another CYP3A gene, CYP3A3, was thought to exist; however, it is now thought that this sequence represents a transcript variant of CYP3A4. Alternatively spliced transcript variants encoding different isoforms have been identified.
Immunogen	Purified recombinant fragment of human CYP3A4 expressed in E. Coli.
Formulation	Purified antibody in PBS with 0.05% sodium azide

Additional Information

Gene ID	1576
Other Names	Cytochrome P450 3A4, 1.14.13.-, 1, 8-cineole 2-exo-monooxygenase, 1.14.13.157, Albendazole monooxygenase, 1.14.13.32, Albendazole sulfoxidase, CYP3A3, CYP3A4, Cytochrome P450 3A3, Cytochrome P450 HLP, Cytochrome P450 NF-25, Cytochrome P450-PCN1, Nifedipine oxidase, Quinine 3-monooxygenase, 1.14.13.67, Taurochenodeoxycholate 6-alpha-hydroxylase, 1.14.13.97, CYP3A4, CYP3A3
Dilution	WB~~1/500 - 1/2000 IHC~~1/200 - 1/1000 FC~~1/200 - 1/400 ICC~~N/A E~~1/10000
Storage	Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CYP3A4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name

CYP3A4 {ECO:0000303|PubMed:11470997,
ECO:0000312|HGNC:HGNC:2637}

Function

A cytochrome P450 monooxygenase involved in the metabolism of sterols, steroid hormones, retinoids and fatty acids (PubMed:[10681376](#), PubMed:[11093772](#), PubMed:[11555828](#), PubMed:[12865317](#), PubMed:[14559847](#), PubMed:[15373842](#), PubMed:[15764715](#), PubMed:[19965576](#), PubMed:[20702771](#), PubMed:[21490593](#), PubMed:[21576599](#)). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase). Catalyzes the hydroxylation of carbon-hydrogen bonds (PubMed:[12865317](#), PubMed:[14559847](#), PubMed:[15373842](#), PubMed:[15764715](#), PubMed:[21490593](#), PubMed:[21576599](#), PubMed:[2732228](#)). Exhibits high catalytic activity for the formation of hydroxyestrogens from estrone (E1) and 17beta- estradiol (E2), namely 2-hydroxy E1 and E2, as well as D-ring hydroxylated E1 and E2 at the C-16 position (PubMed:[11555828](#), PubMed:[12865317](#), PubMed:[14559847](#)). Plays a role in the metabolism of androgens, particularly in oxidative deactivation of testosterone (PubMed:[15373842](#), PubMed:[15764715](#), PubMed:[22773874](#), PubMed:[2732228](#)). Metabolizes testosterone to less biologically active 2beta- and 6beta- hydroxytestosterones (PubMed:[15373842](#), PubMed:[15764715](#), PubMed:[2732228](#)). Contributes to the formation of hydroxycholesterols (oxysterols), particularly A-ring hydroxylated cholesterol at the C- 4beta position, and side chain hydroxylated cholesterol at the C-25 position, likely contributing to cholesterol degradation and bile acid biosynthesis (PubMed:[21576599](#)). Catalyzes bisallylic hydroxylation of polyunsaturated fatty acids (PUFA) (PubMed:[9435160](#)). Catalyzes the epoxidation of double bonds of PUFA with a preference for the last double bond (PubMed:[19965576](#)). Metabolizes endocannabinoid arachidonylethanolamide (anandamide) to 8,9-, 11,12-, and 14,15- epoxyeicosatrienoic acid ethanolamides (EpETE-EAs), potentially modulating endocannabinoid system signaling (PubMed:[20702771](#)). Plays a role in the metabolism of retinoids. Displays high catalytic activity for oxidation of all-trans-retinol to all-trans-retinal, a rate- limiting step for the biosynthesis of all-trans-retinoic acid (atRA) (PubMed:[10681376](#)). Further metabolizes atRA toward 4-hydroxyretinoate and may play a role in hepatic atRA clearance (PubMed:[11093772](#)). Responsible for oxidative metabolism of xenobiotics. Acts as a 2-exo- monooxygenase for plant lipid 1,8-cineole (eucalyptol) (PubMed:[11159812](#)). Metabolizes the majority of the administered drugs. Catalyzes sulfoxidation of the anthelmintics albendazole and fenbendazole (PubMed:[10759686](#)). Hydroxylates antimalarial drug quinine (PubMed:[8968357](#)). Acts as a 1,4-cineole 2-exo-monooxygenase (PubMed:[11695850](#)). Also involved in vitamin D catabolism and calcium homeostasis. Catalyzes the inactivation of the active hormone calcitriol (1-alpha,25-dihydroxyvitamin D(3)) (PubMed:[29461981](#)).

Cellular Location

Endoplasmic reticulum membrane; Single-pass membrane protein.
Microsome membrane; Single-pass membrane protein

Tissue Location

Expressed in prostate and liver. According to some authors, it is not

expressed in brain (PubMed:19094056). According to others, weak levels of expression are measured in some brain locations (PubMed:18545703, PubMed:19359404). Also expressed in epithelium of the small intestine and large intestine, bile duct, nasal mucosa, kidney, adrenal cortex, epithelium of the gastric mucosa with intestinal metaplasia, gallbladder, intercalated ducts of the pancreas, chief cells of the parathyroid and the corpus luteum of the ovary (at protein level).

References

Drug Metab Dispos. 2009 Dec;37(12):2305-13. Biochem Pharmacol. 2010 Jan 15;79(2):277-87.

Images




Figure 1: Western blot analysis using CYP3A4 mAb against human CYP3A4 (AA: 243-430) recombinant protein. (Expected MW is 47.5 kDa)




Figure 2: Western blot analysis using CYP3A4 mAb against HEK293 (1) and CYP3A4 (AA: 243-430)-hIgGFc transfected HEK293 (2) cell lysate.




Figure 3: Immunohistochemical analysis of paraffin-embedded liver cancer tissues using CYP3A4 mouse mAb with DAB staining.




Figure 4: Immunohistochemical analysis of paraffin-embedded human brain tissues using CYP3A4 mouse mAb with DAB staining.




Figure 5: Immunofluorescence analysis of HepG2 cells using CYP3A4 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 6: Flow cytometric analysis of HepG2 cells using CYP3A4 mouse mAb (green) and negative control (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.